

Exercise minimizes weight regain by reducing appetite and burning fat first, carbs later

September 2 2009

Exercise helps prevent weight regain after dieting by reducing appetite and by burning fat before burning carbohydrates, according to a new study with rats. Burning fat first and storing carbohydrates for use later in the day slows weight regain and may minimize overeating by signaling a feeling of fullness to the brain.

The University of Colorado Denver study also found that [exercise](#) prevents the increase in the number of [fat cells](#) that occurs during weight regain, challenging the conventional wisdom that the number of fat cells is set and cannot be altered by dietary or lifestyle changes.

These coordinated physiological changes in the brain and the body lower the 'defended' weight, that is, the weight that our physiology drives us to achieve, and suggest that the effects of exercise on these physiological processes may make it easier to stay on a diet.

The study is "Regular exercise attenuates the metabolic drive to regain weight after long term weight loss." Paul S. MacLean, Janine A. Higgins, Holly R. Wyatt, Edward L. Melanson, Ginger C. Johnson, Matthew R. Jackman, Erin D. Giles, Ian E. Brown and James O. Hill, all of the University of Colorado Denver, conducted the study. The American Physiological Society published the research in the *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*.

How exercise works

Weight gain is, on the surface, remarkably simple, occurring when the calories consumed exceeds the calories expended. On closer examination, the process is remarkably complex. Laboratory animals eat according to physiological signals that may suppress appetite or arouse the desire to eat. These signals are relatively weak in humans, as their intake is largely influenced by psychological, cognitive and lifestyle factors. After dieting, however, the physiological signals emerge to play a more substantial role in controlling intake. Being persistently hungry after losing weight with restricted diets is a big part of the weight regain problem. Most people are unable to ignore this physiological cue and are pushed by their biology to overeat and regain the weight they worked so hard to lose.

Some people are successful at keeping the weight off, and those tracked by The National Weight Control Registry share a number of common characteristics, including a program of regular exercise. The aim of this investigation was to uncover how exercise affects the body's physiology to minimize weight regain.

The researchers used obesity-prone rats. For the first 16 weeks, the rats ate a high-fat diet, as much as they wanted, and remained sedentary. They were then placed on a diet. For the following two weeks, the animals ate a low-fat and low-calorie diet, losing about 14% of their body weight. The rats maintained the weight loss by dieting for eight more weeks. Half the rats exercised regularly on a treadmill during this period while the other half remained sedentary.

In the final 8-weeks, the relapse phase of the study, the rats stopped dieting and ate as much low-fat food as they wanted. The rats in the exercise group continued to exercise and the sedentary rats remained sedentary.

Compared to the sedentary rats, the exercisers:

- regained less weight during the relapse period
- developed a lower 'defended' body weight
- burned more fat early in the day, and more carbohydrates later in the day
- accumulated fewer fat cells and less abdominal fat during relapse
- reduced the drive to overeat
- enhanced the ability to balance energy intake with energy expended

During feeding, the sedentary group preferentially burned carbohydrates while sending fat from the diet to fat tissue. This preferential fuel use stores more calories because it requires less energy to store fat than to store carbohydrates. In addition, burning away the body's carbohydrates may contribute to the persistent feeling of hunger and large appetite of the sedentary animals.

Exercise blunted this fuel preference, favoring the burning of fat for energy needs and saving ingested carbohydrates so that they could be used later in the day. Taken together, the exercise led to a much lower appetite and fewer calories ending up in fat tissue.

The researchers also found that exercise prevented the increase in the number of fat cells observed with weight regain in sedentary rats. In sedentary rats, a population of very small, presumably new, fat cells appears early in the relapse process. Small, new fat cells would not only

accelerate the process of regain, but also increase fat storage capacity in the abdomen. It would also explain why sedentary rats overshoot their previous weight when they relapse.

Conventional wisdom holds that the number of fat cells is determined by genetics, rather than being regulated by diet or lifestyle. Because this effect of exercise is a novel finding, the team will do further research to demonstrate that exercise is, indeed, preventing the formation of new fat cells early in relapse and not simply altering the size of pre-existing fat cells.

More information: ajpregu.physiology.org/cgi/reprint/297/3/R793

Source: American Physiological Society ([news](#) : [web](#))

Citation: Exercise minimizes weight regain by reducing appetite and burning fat first, carbs later (2009, September 2) retrieved 5 May 2024 from <https://medicalxpress.com/news/2009-09-minimizes-weight-regain-appetite-fat.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--