

# Pre-hospital organization: The first links in the chain of survival for heart attack patients

September 1 2009

---

Mortality rate following a heart attack has fallen by more than 50% in Europe over the past 25 years. However, because only minor advances in the medical treatment of AMI are expected over the next decade, it is through organisational changes in the pre-hospital phase that mortality rate will continue this decline to below 5%.

We estimate that acute coronary syndromes (ACS) account for 1.5 million hospital admissions throughout Europe each year. Almost half these patients present with ST-elevation myocardial infarction (STEMI), which are major and immediately life-threatening events.

Just 30 years ago, mortality of acute [myocardial infarction](#) (AMI, [heart attack](#)) in Europe was over 30%. This rate has now dropped dramatically to around 10% within the first month. In clinical trials - where the sickest patients are often excluded - mortality rate is as low as 5%.

This dramatic improvement was initially brought about by the opening of dedicated coronary care units in hospitals. This was followed by 20 years of drug development, and a significant improvement in survival rates. Among the important drugs introduced were those preventing blood from clotting, or even dissolving the clot responsible for blocking the coronary vessel (and so causing the AMI. Other drugs groups include beta-blockers, ACE inhibitors, and statins developed to lower [cholesterol](#) levels. More recently, the treatment of large heart attacks with balloon angioplasty has been a major advance. Although new and better drugs are being developed and drug combinations being refined,

there is less belief in major drug breakthroughs in the next decade.

The pre-hospital phase of AMI treatment has also undergone several changes over the past decade: diagnosis, supported by telemedicine, has improved, and many interventions have been moved from the hospital to the field. It is in this early phase that we must now adopt new collaborations and organisations if mortality rate in this large patient population is to continue its decline. We must adopt new lean principles in the entire organisation of the pre-hospital phase, starting with public awareness of symptoms, and how to raise the alarm. For those patients whose first symptom is cardiac arrest, basic bystander resuscitation should become standard. Despite an abundance of automated external defibrillators (AED) in many regions, their localisation and use are often not well organised. Heart attack victims should call an emergency number, instead of being self or family-transported to the hospital.

There are financial and political disincentives for the transfer of STEMI patients for balloon angioplasty: all of these factors should be addressed:

#### Primary Hospital

- Loss of revenues

#### Referring doctors

- Loss of patients and subsequent revenue
- Pride and unwillingness to admit that another physician can provide better medical care
- Medico-legal liability during transfer

## Lack of organised emergency services

- Private company vs. fire department
- Conflict between firemen and paramedics
- Time constraints: prolonged run times, ECG triage
- What incentive (transfer between hospitals is low paid in some countries)?

The mortality rate following heart attack is today historically low. Further reduction is unlikely to result from new drugs, but there is huge potential in the a leaner organisation of the entire pre-hospital phase. The successful full implementation of these measures has the potential to provide more than 75% reperfusion rate and could reduce mortality to below 5% and approaching the life expectancy rate of the age-matched general population.

Source: European Society of Cardiology ([news](#) : [web](#))

Citation: Pre-hospital organization: The first links in the chain of survival for heart attack patients (2009, September 1) retrieved 23 April 2024 from <https://medicalxpress.com/news/2009-09-pre-hospital-links-chain-survival-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.