

Evaluation of standard liver volume formula for Chinese adults

September 16 2009

Living donor liver transplantation (LDLT) has been used to alleviate the shortage of available liver donors. Accurate estimation of the standard liver volume (SLV) of the living donor and recipient is crucial. Scholars of different countries established several SLV formulae from clinical data, but which formula was the most accurate for Chinese adults has remained unclear.

A research article to be published on August 28, 2009 in the [World Journal of Gastroenterology](#), addresses this question. The research team led by Prof. Lu-Nan Yan from West [China](#) Hospital, Sichuan University evaluated reported SLV formulae using data of 76 cases of LDLT from January 2008 to April 2009.

The results demonstrated that the Chengdu formula was reliable by its applicability in LDLT. Statistical analysis showed that the Fan formula tended to underestimate SLV. The Heinemann, Urata, Vauthey, and Lee formulae overestimated SLV.

With national multicenter data in the future, the Chengdu formula for SLV can be improved. That may be applied to the evaluation of Chinese donors and recipients for LDLT.

More information: Shi ZR, Yan LN, Li B, Wen TF. Evaluation of standard [liver](#) volume formulae for Chinese adults. World J Gastroenterol 2009; 15(32): 4062-4066, www.wjgnet.com/1007-9327/15/4062.asp

Source: World Journal of Gastroenterology ([news](#) : [web](#))

Citation: Evaluation of standard liver volume formula for Chinese adults (2009, September 16)
retrieved 6 May 2024 from

<https://medicalxpress.com/news/2009-09-standard-liver-volume-formula-chinese.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.