

# Teen attitudes toward smoking linked to likelihood of drinking and using drugs

September 30 2009

---

New research by Weill Cornell Medical College researchers looks at the specific ways parents and peers influence teenagers to smoke, drink and use marijuana in combination. Among their findings: attitudes toward smoking influenced teenagers' use of multiple drugs (smoking, drinking and marijuana), and that this manifested itself differently in boys and girls.

For girls, friends were shown to be central. Ambivalent or permissive attitudes within their social group toward smoking were associated with poly-drug use -- defined as two or more of the following behaviors: smoking, [drinking](#) and marijuana use. This wasn't the case with boys, whose poly-drug use was instead predicted by the extent to which they perceived smoking to be prevalent in their larger age group -- not just among their friends.

"If a [teenager](#) feels smoking is socially acceptable and widely practiced, they are much more likely not only to smoke, but to also drink and possibly use marijuana," says lead author Dr. Jennifer A. Epstein, assistant professor of public health in the Division of Prevention and [Health Behavior](#) at Weill Cornell Medical College. "While the differences between how boys and girls are influenced by these social factors are subtle, they could help us develop new gender-specific educational tactics for preventing these behaviors."

The study also revealed several factors that were the same for boys and girls. When their friends drank alcohol or smoked or when their parents

had permissive or ambivalent attitudes toward drinking, both teenage boys and girls were more likely to report poly-drug use. Other major variables included teenagers' inability to refuse drugs and achieve goals through their own efforts.

"A parent's opinion matters. Moms and dads are critical role models and should let their attitudes against drug use be known. It's also important to keep an eye on their child's social circle, since, especially for girls, it's their friends who are so central to influencing their behavior," says Dr. Epstein. "At the same time, parents can do things that reduce their child's risk for using drugs, such as teaching them to set goals and assert themselves."

Researchers analyzed confidential surveys taken by 2,400 sixth- and seventh-graders in inner-city schools in New York City. Questions dealt with substance use and several psychological factors that previous research suggests may be related to drug use. The majority of the schools serve youths from families with incomes averaged well below the federal poverty level.

The current study is one of the first to look at the relationships between [smoking](#), drinking and marijuana use. The vast majority of research in this area has focused on a single substance in isolation, especially among white middle-class suburban populations. The importance of Dr. Epstein's approach is backed up by evidence suggesting that teenage poly-drug use is a significant risk factor for adult poly-drug use.

One implication of these findings, according to Dr. Epstein, is that "comprehensive prevention programs focusing on multiple gateway drugs (alcohol, cigarettes and [marijuana](#)) may prove to be more valuable than programs focusing on a single drug."

More information: The study appears online in the July 1 issue of the

*Journal of Child and Adolescent Substance Abuse.*

Source: New York- Presbyterian Hospital ([news](#) : [web](#))

Citation: Teen attitudes toward smoking linked to likelihood of drinking and using drugs (2009, September 30) retrieved 20 March 2024 from <https://medicalxpress.com/news/2009-09-teen-attitudes-linked-likelihood-drugs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.