

Treatment of personality disorders by psychotherapy: A French multicenter study

September 23 2009

A French multicenter study headed by Jean Cottraux (Lyon) has investigated the role of psychotherapy in borderline personality disorder.

Borderline personality disorder (BPD) is a serious mental illness characterized by pervasive instability in moods, interpersonal relationships, self-image, and behavior. This instability often disrupts family and work life, long-term planning, and the individual's sense of self-identity. Originally thought to be at the "borderline" of psychosis, people with BPD suffer from a disorder of [emotion regulation](#).

To date, there have been no studies comparing cognitive therapy (CT) with Rogerian supportive therapy (RST) in borderline personality disorder. In this study, thirty-three patients were randomly allocated to CT and 32 to RST. The therapists were the same in both groups. Both treatments shared the same duration (1 year) and amount of therapy.

Assessment by independent evaluators utilised the Clinical Global Impression (CGI) Scale, the Hamilton [Depression](#) Scale, Beck Depression Inventory, Beck Anxiety Inventory, Hopelessness Scale, Young Schema Questionnaire II, Eysenck Impulsivity Venturesomeness Empathy (IVE) Inventory, a self-harming behaviours checklist and scales measuring quality of life and the therapeutic relationship. The response criterion was a score of 3 or less on the CGI, associated with a Hopelessness Scale score of

Citation: Treatment of personality disorders by psychotherapy: A French multicenter study (2009, September 23) retrieved 28 April 2024 from <https://medicalxpress.com/news/2009-09-treatment-personality-disorders-psychotherapy-french.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.