

Ulcerative colitis treatment reduces need for surgery by almost half

October 1 2009

A new study led by Mayo Clinic researchers has found that ulcerative colitis patients had a 41 percent reduction in colectomy after a year when treated with infliximab, according to a study published in the October 2009 issue of *Gastroenterology*.

Ulcerative colitis, an [inflammatory bowel disease](#) (IBD) that causes [chronic inflammation](#) of the colon, is characterized by [abdominal pain](#) and diarrhea. Like Crohn's disease, another common IBD, ulcerative colitis can be debilitating and often lead to colectomy or surgical removal of the colon.

"Our purpose in this study was to see if the use of [infliximab](#) for ulcerative colitis would reduce the need for surgery," says William Sandborn, M.D., a Mayo Clinic gastroenterologist and lead author of the study. "We found that treatment with infliximab reduced the need for colectomy by 41 percent compared to patients treated with placebo."

In this multi-center, international study, 728 patients received placebo or infliximab (5 or 10 mg/kg) for 46 weeks and were monitored for hospitalization or surgical outcomes. Eighty-seven percent (630 of 728) had complete follow-up for the endpoint of whether or not they had colectomy, while the remaining 13 percent (98 of 728) of patients had follow-up for less than a year, with a median follow-up of 6.2 months in these patients. The research showed that treatment with infliximab at 0, 2 and 6 and then every 8 weeks reduced the incidence of colectomy through 54 weeks by 41 percent in outpatients with moderately-to-severe

active ulcerative colitis.

The cumulative incidence of colectomy through 54 weeks was 10% for infliximab and 17% for placebo ($p=0.02$). Compared with placebo, fewer ulcerative colitis-related hospitalizations and surgeries/procedures occurred with infliximab therapy.

"One of the most feared outcomes for [ulcerative colitis](#) patients is surgical removal of the colon," says Dr. Sandborn. "Our research hopes to provide other treatment solutions for patients beyond surgery."

Previous research has shown that infliximab therapy induced clinical remission and bowel healing for colitis patients. This new research provides more information and options for patients struggling with this difficult disease, explains Dr. Sandborn.

Source: Mayo Clinic ([news](#) : [web](#))

Citation: Ulcerative colitis treatment reduces need for surgery by almost half (2009, October 1) retrieved 17 April 2024 from <https://medicalxpress.com/news/2009-10-ulcerative-colitis-treatment-surgery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--