

First use of antibody and stem cell transplantation to successfully treat advanced leukemia

November 5 2009

For the first time, researchers at Fred Hutchinson Cancer Research Center have reported the use of a radiolabeled antibody to deliver targeted doses of radiation, followed by a stem cell transplant, to successfully treat a group of leukemia and pre-leukemia patients for whom there previously had been no other curative treatment options.

All fifty-eight patients, with a median age of 63 and all with advanced [acute myeloid leukemia](#) or high-risk myelodysplastic syndrome - a pre-leukemic condition - saw their blood cancers go into remission using a novel combination of low-intensity chemotherapy, targeted radiation delivery by an antibody and a stem-cell transplant. Forty percent of the patients were alive a year after treatment and approximately 35 percent had survived three years, about the same rates as patients who received similar treatment but whose disease was already in remission and who had much more favorable risk for relapse when therapy began.

Results of the research appear online in the journal *Blood*. The principal investigator and corresponding author of the paper is John Pagel, M.D., Ph.D, a transplant oncologist and assistant member of the Hutchinson Center's Clinical Research Division.

The purpose of the study was to find the maximum dose of radiation that patients could tolerate with acceptable [toxic side effects](#), not to assess how effective the novel treatment was, according to Pagel and

colleagues. However, "the results appear to be very encouraging and warrant us to study it further for patients who really have no significant other curative options," Pagel said.

Older (over age 50) patients with active, advanced [leukemia](#) and myelodysplastic syndrome pose the most difficult treatment challenges because standard transplant therapy rarely works, according to Pagel. Both standard and low-dose therapies (a process sometimes known as a "mini transplant" and pioneered at the Hutchinson Center) used to kill [leukemia cells](#) in the bloodstream in preparation for a transplant usually require that patients be in remission.

The patients in this study, who came from all over the world to participate in the Phase 1 clinical trial, were in large part those with active relapsed disease that in many cases had failed to respond to standard therapies. Eighty-six percent of the 58 patients had active disease and only 10 percent were in remission when therapy was begun. Their cancers had failed previous treatment attempts. "These were people who had extremely advanced high-risk disease, they were typically older - most of them were in their 60s and some were in their 70s - and had few or no other options for a potential cure. In fact most, if not all, would not been offered a stem cell transplant here or elsewhere. It is fair to say that these patients would likely have died without a transplant being performed if they had not been given the opportunity to participate in this study."

To find the optimal dose of radiation, researchers began at 12 Gy (Gray, a unit of measurement of absorbed [radiation dose](#)) and escalated the dosages in increments of 2 Gy up to a Gy of 26. At that dose, some toxicity to the heart and lungs was found so they concluded 24 Gy to be the maximum effective dosage. The 21 patients who received the maximum radiation dose have survived the longest, researchers reported.

The key to success in this study was use of a radiolabeled antibody that has therapeutic iodine 131 attached and is designed to target leukemic blood cells that carry a marker on the surface of the cell known as CD45. Its use in delivering targeted amounts of radiation was developed several years ago at the Hutchinson Center. Delivered intravenously, the radiation looks for the CD45 antigen receptor on the surface of blood cells. This approach results in a two- to four-fold increase in the amount of radiation that reaches cancerous cells as compared to standard external beam radiation, which also radiates normal surrounding organs and tissue. The more radiation that can be applied, the more cancer cells will be killed in preparation for donor stem cells to take over the diseased immune system and kill off the remaining cancer cells.

Pagel said further research is needed to test more patients at the highest radiation dose both at the Hutchinson Center and at other transplant centers around the country.

Source: Fred Hutchinson Cancer Research Center ([news](#) : [web](#))

Citation: First use of antibody and stem cell transplantation to successfully treat advanced leukemia (2009, November 5) retrieved 3 May 2024 from <https://medicalxpress.com/news/2009-11-antibody-stem-cell-transplantation-successfully.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--