

Burned out, depressed surgeons more likely to commit more major medical errors

November 23 2009

Surgeons who are burned out or depressed are more likely to say they had recently committed a major error on the job, according to the largest study to date on physician burnout. The new findings suggest that the mental well-being of the surgeon is associated with a higher rate of self-reported medical errors, something that may undermine patient safety more than the fatigue that is often blamed for many of the medical mistakes.

Although [surgeons](#) do not appear more likely to make mistakes than physicians in other disciplines, surgical errors may have more severe consequences for patients due to the interventional nature of the work. Some estimate that as many as 10 percent of hospitalized patients are impacted by medical errors.

"People have talked about [fatigue](#) and long working hours, but our results indicate that the dominant contributors to self-reported medical errors are [burnout](#) and depression," said Charles M. Balch, M.D., a professor of surgery at the Johns Hopkins University School of Medicine and one of the study's leaders. "All of us need to take this into account to a greater degree than in the past. Frankly, burnout and depression hadn't been on everybody's radar screen."

Nine percent of the 7,905 surgeons who responded to a June 2008 survey commissioned by the American College of Surgeons for a study led by researchers from Johns Hopkins University School of Medicine and the Mayo Clinic reported having made a major medical mistake in the

previous three months. Overall, 40 percent of the surgeons who responded to the survey said they were burned out.

Researchers asked a variety of questions, including queries that rated three elements of burnout —emotional exhaustion, depersonalization and personal accomplishment — and others that screened for depression.

Each one-point increase on a scale that measured depersonalization — a feeling of withdrawal or of treating patients as objects rather than as human beings — was associated with an 11 percent increase in the likelihood of reporting an error. Each one-point increase on a scale measuring emotional exhaustion was associated with a 5 percent increase.

Mistakes also varied by specialty. Surgeons practicing obstetrics/gynecology and plastic surgery were much less likely to report errors than general surgeons.

Researchers acknowledged the limitations of self-reporting surveys, saying they couldn't tell from their research whether burnout and depression led to more medical errors or whether [medical errors](#) triggered burnout and depression among the surgeons who made the mistakes.

The results are being published online today in the *Annals of Surgery* and will be published in the printed journal in an upcoming issue.

Notably, the research shows that the number of nights on call per week and the number of hours worked were not associated with reported errors after controlling for other factors.

"The most important thing for those of us who work with other surgeons who do not appear well is to address it with them so that they can get the

help they need," says Julie A. Freischlag, M.D., chair of the Department of Surgery at the Johns Hopkins University School of Medicine and another of the study's authors.

Source: Johns Hopkins Medical Institutions

Citation: Burned out, depressed surgeons more likely to commit more major medical errors (2009, November 23) retrieved 25 April 2024 from <https://medicalxpress.com/news/2009-11-depressed-surgeons-commit-major-medical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.