

# Factors from common human bacteria may trigger multiple sclerosis

November 24 2009

---

Current research suggests that a common oral bacterium may exacerbate autoimmune disease. The related report by Nichols et al, "Unique Lipids from a Common Human Bacterium Represent a New Class of TLR2 Ligands Capable of Enhancing Autoimmunity," appears in the December 2009 issue of *The American Journal of Pathology*.

Multiple sclerosis (MS), a disease where the [immune system](#) attacks the brain and spinal cord, affects nearly 1 in 700 people in the United States. Patients with multiple sclerosis have a variety of neurological symptoms, including muscle weakness, difficulty in moving, and difficulty in speech.

*Porphyromonas gingivalis*, a common oral bacterium in humans, produces a unique type of [lipid](#), phosphorylated dihydroceramides (DHCs), which enhance inflammatory responses. These lipids are also likely produced by bacteria found in other parts of the body including the [gastrointestinal tract](#). To determine if these lipids accentuate immune-mediated damage in autoimmune disease, researchers led by Robert B. Clark and Frank C. Nichols of the University of Connecticut Health Center administered phosphorylated DHCs in a mouse model of MS. The severity of disease was significantly enhanced by the addition of these lipids in a manner that was dependent on activation of the immune system. These data suggest that phosphorylated DHCs from bacteria commonly found in humans may trigger or increase the severity of [autoimmune diseases](#) such as multiple sclerosis.

The authors state that "while it is clear that the immune system in most individuals has the potential to attack self-tissues, the "tipping" factors that initiate and propagate autoimmune diseases such as [multiple sclerosis](#) in only a subset of individuals remain unknown. Overall, [their] results represent the first description that phosphorylated DHCs derived from common human bacteria are capable of enhancing autoimmune disease." Thus, these lipids may function as "tipping" factors, playing a previously unrecognized role in initiating or exacerbating human autoimmune diseases. In future studies, Dr. Clark and colleagues plan to characterize the effects of phosphorylated DHCs on specific cells of the immune system and to identify how and where these lipids are deposited in tissues throughout the body. In addition to the role of these lipids in triggering and worsening MS, the authors believe that phosphorylated DHCs may have the potential to serve both as new markers of MS disease activity and as new targets for therapeutic intervention.

More information: Nichols FC, Housley W, O'Connor C, Manning T, Wu S, Clark RB: Unique Lipids from a Common Human Bacterium Represent a New Class of TLR2 Ligands Capable of Enhancing Autoimmunity. *Am J Pathol* 175: 2430-2438

Source: [American Journal of Pathology](#) ([news](#) : [web](#))

Citation: Factors from common human bacteria may trigger multiple sclerosis (2009, November 24) retrieved 6 May 2024 from <https://medicalxpress.com/news/2009-11-factors-common-human-bacteria-trigger.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---