

## Theory about long and short-term memory questioned

November 9 2009

The long-held theory that our brains use different mechanisms for forming long-term and short-term memories has been challenged by new research from UCL, published today in *PNAS*.

Neuroscientists formed this theory based on observation of patients with amnesia, a condition that severely disrupts the ability to form longlasting memories. Typically, amnesia is caused by injury to the hippocampi, a pair of brain structures located in the depth of the temporal lobes.

Despite the condition devastating long-term <u>memory</u>, such patients are quite proficient in rehearsing a phone number over short periods of time, as long as their attention is not distracted. This led to a hypothesis that the <u>hippocampus</u> supports long-term but not short-term memory. However, the UCL study shows that this distinction now needs to be reconsidered.

The team studied patients with a specific form of <u>epilepsy</u> called 'temporal lobe epilepsy with bilateral hippocampal sclerosis', which leads to marked dysfunction of the hippocampi. They asked the patients to try and memorise photographic images depicting normal scenes, for example chairs and a table in a living-room. Their memory of the image was tested and brain activity recorded using MEG (<u>magnetoencephalography</u>) after a short interval of just five seconds, or a long interval of 60 minutes.



The researchers discovered that short-term memory about more detailed aspects of the scene, for example whether the table was located left or right of the chairs, required the coordinated activity of a network of visual and temporal brain areas, whereas standard short-term memory drew on a very different network. Critically, the coordinated activity of visual and temporal brain areas was disrupted in the patients with hippocampal sclerosis.

Professor Emrah Duzel, UCL Institute for <u>Cognitive Neuroscience</u>, said: "As we anticipated, the patients could not distinguish the studied images from new images after 60 minutes - but performed normally at five seconds. However, a striking deficit emerged even at five seconds when we asked them to recall the detailed arrangement of objects within the scenes.

"These findings identify two distinct short-term memory networks in the brain: one that functions independently of the hippocampus and remains intact in patients with long-term memory deficits and one that is dependent on the hippocampus and is impaired alongside long-term memory."

Nathan Cashdollar, UCL Institute of Neurology and first author of the paper, added: "Recent behavioural observations had already begun challenging the classical distinction between long-term and short-term memory which has persisted for nearly half a century. However, this is the first functional and anatomical evidence showing which mechanisms are shared between short-term and long-term memory and which are independent."

"They also highlight that patients with impaired long-term memory have a short-term memory burden to carry in their daily life as well."

More information: Cashdollar N., Malecki U., Rugg-Gunn F.J., Duncan



J.S., Lavie N., Duzel E. 'Hippocampus dependent and independent thetanetworks of active maintenance' is published in the *Proceedings of the National Academy of Sciences* of the United States of America (*PNAS*) in this week's online early edition.

Source: University College London (<u>news</u> : <u>web</u>)

Citation: Theory about long and short-term memory questioned (2009, November 9) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2009-11-theory-short-term-memory.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.