

National survey tracks rates of common mental disorders among American youth

December 14 2009

Only about half of American children and teenagers who have certain mental disorders receive professional services, according to a nationally representative survey funded in part by the National Institute of Mental Health (NIMH). The survey also provides a comprehensive look at the prevalence of common mental disorders.

The results are part of the National Health and Nutrition Examination Survey (NHANES), a collaboration between NIMH and the National Center for Health Statistics of the Centers for Disease Control and Prevention. The survey conducted from 2001 to 2004 had 3,042 participants. These most recent results include data from children and adolescents ages 8 to 15, and were published online ahead of print December 14, 2009, in the journal *Pediatrics*.

"Data on the prevalence of mental disorders among U.S. youth have been varied, making it difficult to truly understand how many children and teens are affected," said NIMH Director Thomas R. Insel, M.D. "These data from the NHANES survey can serve as an important baseline as we follow trends of mental disorders in children."

In the study, the young people were interviewed directly. Parents or caregivers also provided information about their children's mental health. The researchers tracked six mental disorders—generalized anxiety disorder (GAD), panic disorder, eating disorders (anorexia and bulimia), depression, [attention deficit hyperactivity disorder \(ADHD\)](#) and conduct disorder. The participants were also asked about what

treatment, if any, they were receiving.

Overall, 13 percent of respondents met criteria for having at least one of the six mental disorders within the last year. About 1.8 percent of the respondents had more than one disorder, usually a combination of ADHD and conduct disorder. Among the specific disorders,

- 8.6 percent had ADHD, with males more likely than females to have the disorder;
- 3.7 percent had depression, with females more likely than males to have the disorder;
- 2.1 percent had conduct disorder;
- 0.7 percent had an anxiety disorder (GAD or panic disorder);
- 0.1 percent had an eating disorder (anorexia or bulimia).

"With the exception of ADHD, the prevalence rates reported here are generally lower than those reported in other published findings of mental disorders in children, but they are comparable to other studies that employed similar methods and criteria," said lead author Kathleen Merikangas, Ph.D., of NIMH.

Those of a lower socioeconomic status were more likely to report any disorder, particularly ADHD, while those of a higher socioeconomic status were more likely to report having an anxiety disorder. Mexican-Americans had significantly higher rates of mood disorders than whites or African-Americans, but overall, few ethnic differences in rates of disorders emerged.

Merikangas and colleagues also found that overall, 55 percent of those with a disorder had consulted with a mental health professional, confirming the trend of an increase in service use for childhood mental disorders, especially ADHD. However, only 32 percent of youth with an anxiety disorder sought treatment, a finding consistent with other studies. Moreover, African-Americans and Mexican-Americans were significantly less likely to seek treatment than whites, reiterating the need to identify and remove barriers to treatment for minority youth, noted the researchers.

"Until now, there has been a dearth of reliable data on the magnitude, course and treatment patterns of mental disorders among U.S. youth," said Dr. Merikangas. "When combined with data from other nationally representative surveys, the data will provide a valuable basis for making decisions about health care for American youth," she concluded.

Source: NIH/National Institute of Mental Health

Citation: National survey tracks rates of common mental disorders among American youth (2009, December 14) retrieved 23 April 2024 from <https://medicalxpress.com/news/2009-12-national-survey-tracks-common-mental.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--