

# Seeing with your hands? Hand amputation may result in altered perception around the hands

January 6 2010

---

where actions such as grasping and touching occur — is known as the "action space." Research has shown that visual information in this area is organized in hand-centered coordinates — in other words, the representation of objects in the human brain depends on their physical location with respect to the hand. According to new research in *Psychological Science*, a journal of the Association for *Psychological Science*, amputation of the hand results in distorted visuospatial perception (i.e., figuring out where in space objects are located) of the action space.

Scientists Tamar R. Makin, Meytal Wilf, and Ehud Zohary from the Hebrew University of Jerusalem along with Isabella Schwartz from Hadassah Mount Scopus Hospital in Jerusalem wanted to investigate how [hand](#) amputations affect visuospatial [perception](#) in near space.

Volunteers with either left- or right-hand amputations participated in this experiment. Volunteers were instructed to look at a central cross on a screen as two white squares were briefly shown to the left and right side of the cross. The volunteers had to indicate which of the squares was further away from the cross.

The results reveal that hand amputations affect visuospatial perception. When the right square was slightly further away from the center, participants with right-hand amputations tended to perceive it as being at the same distance from the center as the left square; this suggests that

these volunteers underestimated the distance of the right square relative to the left. Conversely, when the left square was further away, participants with left-hand amputations perceived both squares as being equally far away from the center — these participants underestimated the left side of near space. Interestingly, when the volunteers were seated farther away from the screen, they were more accurate in judging the distances, indicating that hand amputations may only affect perception of the space close to the body.

The findings suggest that losing a hand may shrink the action space on the amputated side, leading to permanent distortions in spatial perception. According to the researchers, "This shows that the possibility for action in near space shapes our perception — the space near our hands is really special, and our ability to move in that space affects how we perceive it."

The researchers note that these results have implications for spatial hemineglect — a condition (often following brain injury) in which the patient cannot perceive objects on one side of space. This condition is very often associated with paralysis of the hand in the neglected side, which, according to the current study, might intensify the perceptual neglect. The authors suggest that, based on their findings, "current rehabilitation approaches that emphasize action on the affected side may reverse this process." For example, encouraging the use of the affected hand or by providing visual feedback (through prism adaptation or mirrors) may help overcome hemineglect by increasing the size of the action space on the affected side.

Provided by Association for Psychological Science

Citation: Seeing with your hands? Hand amputation may result in altered perception around the hands (2010, January 6) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2010-01-amputation-result-perception.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.