

Green tea could modify the effect of cigarette smoking on lung cancer risk

January 12 2010

Drinking green tea could modulate the effect of smoking on lung cancer. Results of this hospital-based, randomized study conducted in Taiwan were presented at the AACR-IASLC Joint Conference on Molecular Origins of Lung Cancer.

"[Lung cancer](#) is the leading cause of all cancer deaths in Taiwan," said I-Hsin Lin, M.S., a student at Chung Shan Medical University in Taiwan. "Tea, particularly green tea, has received a great deal of attention because tea polyphenols are strong antioxidants, and tea preparations have shown inhibitory activity against tumorigenesis."

However, previous studies of green tea have been inhibited by the flaws of the epidemiologic model with its inherent biases.

Lin and colleagues enrolled 170 patients with lung cancer and 340 healthy patients as controls. The researchers administered questionnaires to obtain demographic characteristics, cigarette [smoking](#) habits, green tea consumption, dietary intake of fruits and [vegetables](#), cooking practices and family history of lung cancer. They also performed genotyping on insulin-like growth factors as polymorphisms on the following insulin-like growth factors: IGF1, IGF2 and IGFBP3, which have all been reported to be associated with cancer risk.

Among smokers and non-smokers, those who did not drink green tea had a 5.16-fold increased risk of lung cancer compared with those who drank at least one cup of green tea per day. Among smokers, those who did not

drink green tea at all had a 12.71-fold increased risk of lung cancer compared with those who drank at least one cup of green tea per day.

Lin and colleagues suspect genetics may play a role in this risk differential. Green tea drinkers with non-susceptible IGF1 (CA)19/(CA)19 and (CA)19/X genotypes reported a 66 percent reduction in lung cancer risk as compared with green tea drinkers carrying the IGF1 X/X genotype.

Heavy smokers carrying susceptible IGF1, IGF2 and IGFBP3 genotypes also had a higher risk of lung cancer compared with nonsmokers carrying non-susceptible IGF1, IGF2 and IGFBP3 genotypes.

"Our study may represent a clue that in the case of lung cancer, smoking-induced carcinogenesis could be modulated by [green tea](#) consumption and the growth factor environment," said Lin.

Provided by American Association for Cancer Research

Citation: Green tea could modify the effect of cigarette smoking on lung cancer risk (2010, January 12) retrieved 2 May 2024 from <https://medicalxpress.com/news/2010-01-green-tea-effect-cigarette-lung.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--