

Healthy older adults with subjective memory loss may be at increased risk for MCI and dementia

January 8 2010

Forgot where you put your car keys? Having trouble recalling your colleague's name? If so, this may be a symptom of subjective cognitive impairment (SCI), the earliest sign of cognitive decline marked by situations such as when a person recognizes they can't remember a name like they used to or where they recently placed important objects the way they used to. Studies have shown that SCI is experienced by between one-quarter and one-half of the population over the age of 65. A new study, published in the January 11, 2010, issue of the journal *Alzheimer's & Dementia*, finds that healthy older adults reporting SCI are 4.5 times more likely to progress to the more advanced memory-loss stages of mild cognitive impairment (MCI) or dementia than those free of SCI.

The long-term study completed by researchers at NYU Langone Medical Center tracked 213 adults with and without SCI over an average of seven years, with data collection taking nearly two decades. Further cognitive decline to MCI or dementia was observed in 54 percent of SCI persons, while only in 15 percent of persons free of SCI.

"This is the first study to use mild [cognitive impairment](#) as well as dementia as an outcome criterion to demonstrate the outcome of SCI as a possible forerunner of eventual Alzheimer's disease," said Barry Reisberg, MD, professor of psychiatry, director of the Fisher Alzheimer's Disease Program and director, Clinical Core, NYU

Alzheimer's Disease Center at NYU Langone Medical Center. "The findings indicate that a significant percentage of people with early subjective symptoms may experience further [cognitive decline](#), whereas few persons without these symptoms decline. If decline does occur in those without SCI symptoms, it takes considerably longer than for those with subjective cognitive symptoms."

According to the authors, scientists and physicians can now target the prevention of eventual Alzheimer's disease in the SCI stage, beginning more than 20 years before [dementia](#) becomes evident

"These intriguing results more fully describe the possible relationship between early signs of memory loss and development of more serious impairment. This is critical to know, as we look for ways to define who is at risk and for whom the earliest interventions might be successful," said Neil Buckholtz, PhD, National Institute on Aging (NIA) which supported the research. "These findings also underscore the importance of clinicians' asking about, and listening to, concerns regarding changes in cognition and memory among their aging patients."

Provided by New York University School of Medicine

Citation: Healthy older adults with subjective memory loss may be at increased risk for MCI and dementia (2010, January 8) retrieved 6 May 2024 from <https://medicalxpress.com/news/2010-01-healthy-older-adults-subjective-memory.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--