

# Insecticide-treated net program in Tanzania

January 11 2010

---

A Tanzanian program to supply insecticide treated nets to prevent malaria in pregnant women successfully delivers the intervention to only 30% of women, but simple changes could increase effectiveness, according to an article in *CMAJ* (*Canadian Medical Association Journal*).

Insecticide treated nets have been shown to reduce childhood mortality by 17% in different areas in Africa, reduce infection with malaria parasites in [pregnant women](#) by 38%, and decrease the incidence of severe malarial anemia by 47%.

The study by researchers from the London School of Hygiene and Tropical Medicine in London, England and the Ifakara Health Institute, Dar es Salaam, Tanzania looked at a sample of 6198 households that participated in the Tanzanian National Voucher Scheme between 2005 and 2007.

The program targets pregnant women and infants by providing vouchers for nets to women at their first antenatal visit. However, the several simple steps involved in the program result in some attrition at each step, which cumulates in a lower success rate than desired.

The researchers suggest that process analysis is useful to identify how programmes can be improved. Nets pretreated with [insecticide](#) are now being delivered, rather than relying on the recipients to treat the nets.

**More information:** <http://www.cmaj.ca/embargo/cmaj090268.pdf>

Provided by Canadian Medical Association Journal

Citation: Insecticide-treated net program in Tanzania (2010, January 11) retrieved 6 May 2024 from <https://medicalxpress.com/news/2010-01-insecticide-treated-net-tanzania.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.