

Kidney abnormalities require more research

January 14 2010

Abnormalities in the kidneys and their blood vessels occur in at least 25% of healthy individuals, according to a study appearing in an upcoming issue of the *Clinical Journal of the American Society Nephrology* (CJASN). While most of these conditions are not harmful enough to prevent someone from donating a kidney, future studies are needed to determine their impact on long-term health.

When patients undergo computerized tomography (CT) imaging scans for medical reasons, physicians sometimes uncover unexpected abnormal findings in the kidneys or in the arteries that supply blood to the kidneys. To determine how commonly these abnormalities occur in healthy adults who have no clinical illness to justify a [CT scan](#), Elizabeth Lorenz, MD, and Andrew Rule, MD (Mayo Clinic) examined nearly 2000 adults who came in to the Mayo Clinic to see if they could donate a kidney to a patient with [kidney failure](#). These adults all underwent CT scans with detailed imaging of the kidneys and renal arteries.

The investigators found that abnormalities in the kidneys and renal arteries were quite common, occurring in at least 1 in 4 adults. [Kidney stones](#) were the most common abnormality and were present in 1 in 10 adults. Some abnormalities such as narrowing of the renal arteries, scars in the kidney, and masses in the kidney (several of which were early cancers) were found more often in patients over 50 years of age. Other abnormalities were more common in women than men.

The majority of these abnormalities were mild and were not causing the individuals any discomfort. More than 73% of these abnormalities were

not harmful enough to prevent kidney donation, and only rarely did physicians prescribe medications to treat [renal artery](#) abnormalities or to prevent further [kidney](#) stone formation and growth.

"These findings highlight an interesting challenge implicit with improvements in imaging technology: physicians are now finding more subtle abnormalities in the kidneys and renal arteries, but lack clear evidence as to whether these findings are benign or harmful to the long-term health of patients," said Dr. Lorenz.

More information: The article, entitled "Prevalence of Renal Artery and Kidney Abnormalities By Computed Tomography among Healthy Adults," will appear online on January 14, 2009, [doi:10.2215/CJN.07641009](https://doi.org/10.2215/CJN.07641009)

Provided by American Society of Nephrology

Citation: Kidney abnormalities require more research (2010, January 14) retrieved 26 April 2024 from <https://medicalxpress.com/news/2010-01-kidney-abnormalities-require.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--