

Do men with early prostate cancer commit suicide more frequently?

February 25 2010

The introduction of prostate-specific antigen (PSA) testing as a screening tool for early detection of prostate cancer (PCa) in the beginning of the 1990s drastically increased the detection of PCa. The risk of suicide is increased among cancer patients including men with PCa. To assess the risk of suicide among men diagnosed with PCa subsequent to PSA testing, a nation-wide study was carried out in Sweden. The results are published in the March issue of *European Urology*, the scientific journal of the European Association of Urology (EAU).

Anxiety related to a crisis reaction may develop into a depression, and several studies have shown that there is a high [anxiety level](#) among screeners in various screening programs. However, as in most countries, men who underwent PSA testing in Sweden at the time represent an opportunistic screening population and not a true population-based screening program by invitation. Therefore, they may have been more health conscious, less prone to develop depression, and more prepared to accept the potential side effects of curative treatment than the general population.

The number of suicides registered for cases in the [Prostate Cancer](#) Base Sweden (a database in which a number of different registers are merged) cohort was compared with the expected number of suicides in an age-matched general male Swedish population. The strengths of this study include the population-based design with inclusion of approximately 98% of all men in Sweden diagnosed with PCa between 1997 and 2006.

There was no evidence for an increased risk of [suicide](#) among men diagnosed with early nonpalpable PCa detected by PSA testing. The suicide rate, however, was twice as high among men diagnosed with locally advanced or metastatic disease compared with the general male population. This is important to acknowledge in order to focus on the need to identify signs of depression and optimise treatment among this category of patients.

More information: Suicide Risk in Men with Prostate-Specific Antigen-Detected Early Prostate Cancer: A Nationwide Population-Based Cohort Study from PCBaSe Sweden, Anna Bill-Axelsson, Hans Garmo, Mats Lambe, Ola Bratt, Jan Adolfsson, Ullakarin Nyberg, Gunnar Steineck, Pär Stattin. *European Urology* Volume 57, issue 3, pages 363-550, March 2010

Provided by European Association of Urology

Citation: Do men with early prostate cancer commit suicide more frequently? (2010, February 25) retrieved 20 April 2024 from <https://medicalxpress.com/news/2010-02-men-early-prostate-cancer-commit.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.