

Imaging fat layer around heart can help predict disease

March 15 2010

Imaging epicardial adipose tissue, or the layer of fat around the heart, can provide extra information compared with standard diagnostic techniques such as coronary artery calcium scoring, according to research by cardiologists at Emory University School of Medicine. The size of the layer of fat around the heart can be measured by X-ray imaging techniques such as CT or MRI.

"This information may be used as a 'gate keeper', in that it could help a cardiologist decide whether a patient should go on to have a nuclear stress test," says Paolo Raggi, MD, professor of medicine (cardiology) and radiology and director of Emory's <u>cardiac imaging</u> center.

Results from two studies were presented at the American College of Cardiology meeting in Atlanta on Sunday, March 14.

The first study, presented by cardiology fellow Nikolaos Alexopoulos, MD, now at the University of Athens, Greece, shows that patients with a larger volume of epicardial adipose tissue tend to have the types of atherosclerotic plaques cardiologists deem most dangerous: non-calcified plaques.

Calcium tends to build up in <u>atherosclerotic plaques</u>. Even though the heart's overall coronary calcium burden is a good predictor of heart disease, calcium in an individual plaque doesn't necessarily mean imminent trouble, Raggi says. Researchers have been learning that non-calcified plaques indicate active buildup in that coronary artery, and



studies suggest that the <u>fat</u> around the heart secretes more inflammatory hormones, compared to the fat just under the skin.

"Release of inflammatory factors from epicardial adipose tissue may be promoting an active atherosclerotic process, and this is indicated by the presence of non-calcified plaques," Raggi says.

Emory researchers examined 214 patients through <u>cardiac CT</u>, and performed coronary artery scoring as well as assessing the patients' epicardial adipose tissue volume and the plaque in their coronary arteries. The epicardial adipose tissue volume was highest in the patients with non-calcified plaques (roughly 60 percent more than those with calcified plaques).

The second study, presented by Emory cardiology fellow Matthew Janik, MD, measured epicardial fat in patients receiving a nuclear stress test. The 382 patients had chest pain but did not have known cardiovascular disease. A nuclear stress test picks up signs of inducible ischemia: deficiencies in blood flow in the <u>heart</u> muscle.

Here, the researchers found that the presence of ischemia correlated more closely with epicardial <u>adipose tissue</u> volume than with the coronary calcium score.

More information:

References:

N. Alexopoulos et al. Epicardial adipose tissue and coronary artery plaque characteristics. Atherosclerosis, in press (2009).

N. Alexopoulos and P. Raggi. Calcification in atherosclerosis. Nature Reviews Cardiology 6, 681-688 (Nov 2009).



Provided by Emory University

Citation: Imaging fat layer around heart can help predict disease (2010, March 15) retrieved 4 May 2024 from <u>https://medicalxpress.com/news/2010-03-imaging-fat-layer-heart-disease.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.