

Researchers find four biomarkers important in colorectal cancer treatment prognosis

April 19 2010

Researchers at the University of Alabama at Birmingham (UAB) Department of Pathology have discovered a set of four biomarkers that will help predict which patients are more likely to develop aggressive colorectal cancer and which are not. The findings also shed light on the genetics that result in worse colorectal cancer-treatment outcomes for African-Americans, compared with Caucasians, the researchers said.

In data presented April 19 at the American Association of Cancer Research annual meeting in Washington, D.C., Liselle Bovell, a graduate student working in the laboratory of UAB Associate Professor of Pathology Upender Manne, Ph.D., discovered that patients who tested positive for higher levels of a genetic [biomarker](#) called microRNA ([miRNA](#)) had increased risk of death after being treated for colorectal cancer.

In genetic tests of tumor samples from the patients, the presence of higher levels of miRNA-21 and miR-106a signaled poorer prognosis after treatment for both Caucasian and African-American patients compared with patients who did not have the higher miRNA levels. The presence of higher levels of miR-181b and miR-203 signaled poorer prognosis after treatment for African-American patients, but not for Caucasians.

"This knowledge gives us solid, prognostic information, so we can better manage patients with these cancers early after diagnosis or surgery," Bovell said.

Nobel Laureate Philip A. Sharp, Ph.D., of the Koch Institute for Integrative Cancer Research at the Massachusetts Institute of Technology, hosted a press conference at the annual meeting that included Bovell and other researchers. In a statement, Sharp said, "The science of miRNAs and related small RNAs will continue to generate new insights into cancer and possible future treatment."

The UAB researchers analyzed data from 218 patients who underwent surgery and other treatments for colorectal cancer at UAB Hospital between 1982 and 2004. They examined the gene-test results of tumor and benign-tissue samples and discovered the four miRNAs important to assessing patient survival.

For many cancers, including colon cancer, African-Americans have lower survival rates than whites. Possible reasons behind this disparity — including genetic variation, tumor characteristics, access to health care and other factors — are being examined extensively.

"Our findings underscore the potential clinical usefulness of miRNAs in studying cancer risk and [cancer](#) progression, and we've shown that race and ethnicity should be considered in the evaluation," Manne said.

Provided by University of Alabama at Birmingham

Citation: Researchers find four biomarkers important in colorectal cancer treatment prognosis (2010, April 19) retrieved 25 April 2024 from <https://medicalxpress.com/news/2010-04-biomarkers-important-colorectal-cancer-treatment.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.