

Study: Carbon monoxide exposure can be reduced during routine anesthesia in kids

April 29 2010

Doctors at Children's National Medical Center have found that carbon monoxide levels in the blood of young children increase during routine general anesthesia. Anesthesiologists have found for the first time that, under certain circumstances, infants and children may be exposed to carbon monoxide during routine anesthesia resulting in a rise in the carbon monoxide levels in the child's blood.

Because carbon monoxide can be generated as a <u>byproduct</u> of anesthesia, anesthesiologists world-wide use specific precautions to prevent and limit its production, according to Richard J. Levy, MD, Chief of Cardiac Anesthesiology, at Children's National. Dr. Levy's team identified the conditions in which carbon monoxide may be inhaled during anesthesia:

- The first study demonstrated that carbon monoxide detected in the breathing circuit correlated with the increase in blood levels in children 2 years and older. The study included 15 patients between 4 months and 8 years.
- The second study identified that the patient's own exhaled carbon monoxide may be "re-breathed" during low-flow anesthesia the current standard of care where fresh gas flows more slowly into the circuit, rather than rapidly.

"The main goal is to provide the safest environment for young patients who require surgery," said Dr. Levy. "We have identified tangible ways



to reduce the risk of carbon monoxide exposure, and our hope is that these changes will be implemented internationally."

Much remains unknown about the effects of low-dose carbon monoxide exposure on the developing brain. Several recent studies have suggested there may be a link to hearing impairments. Though there is more research to be done to determine these impacts, Dr. Levy recommends two immediate changes that will eliminate the risk of <u>carbon monoxide</u> exposure in children:

- In the anesthesia machine, use <u>carbon dioxide</u> absorbents that lack strong metal alkali and do not degrade inhaled anesthetics (avoids CO production risk)
- Avoid of low-flow anesthesia (avoids CO re-breathing risk)

Following Dr. Levy's study findings, Children's National switched to the recommended absorbent to minimize the risk to patients. Although the sample size is small, the results are compelling and Dr. Levy believes changes should be implemented.

Dr. Levy's research has been named the top study out of 600 at the International Anesthesia Research Society 2010 meeting, and won the John J. Downes Research Award for best abstract at the upcoming Society of Pediatric <u>Anesthesia</u>/American Academy of Pediatrics 2010 Annual meeting.

More information: Read the study: <u>www.anesthesia-analgesia.org/c</u> ... t/110/3/747.abstract



Provided by Children's National Medical Center

Citation: Study: Carbon monoxide exposure can be reduced during routine anesthesia in kids (2010, April 29) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2010-04-carbon-monoxide-exposure-routine-anesthesia.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.