

Endoscopic resection of a large ileal lipoma

April 12 2010

A research team from Japan reported a case of a large lipoma, 50 mm in diameter, in the terminal ileum, which was completely removed by a combination of the so-called endoscopic unroofing technique and endoscopic submucosal dissection.

Lipoma is relatively common in the colon but is less often in the [small intestine](#). Most lipomas are incidentally detected at [endoscopy](#) and are usually small and asymptomatic. However, some of them can present with obstruction and/or intussusceptions. Surgical resection is commonly recommended to remove such significant lipomas with a limited pedicle and larger than 2 cm in size, as endoscopic resection may result in unfavorable complications such as intestinal perforations.

A research article to be published on April 7, 2010 in the [World Journal of Gastroenterology](#) addresses this question. The endoscopic submucosal dissection (ESD) technique was used to remove a large lesion en bloc.

Their results showed large ileal lipoma can be completely removed with the modified ESD procedure in combination with endoscopic unroofing technique (from the top to the bottom like peeling a banana).

More information: Morimoto T, Fu KI, Konuma H, Izumi Y, Matsuyama S, Ogura K, Miyazaki A, Watanabe S. Peeling a giant ileal lipoma with endoscopic unroofing and endoscopic submucosal dissection. www.wjgnet.com/1007-9327/full/v16/i13/1676.htm

Provided by World Journal of Gastroenterology

Citation: Endoscopic resection of a large ileal lipoma (2010, April 12) retrieved 23 April 2024 from <https://medicalxpress.com/news/2010-04-endoscopic-resection-large-ileal-lipoma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.