

Researchers make first direct recording of mirror neurons in human brain

April 12 2010



(PhysOrg.com) -- Mirror neurons, many say, are what make us human. They are the cells in the brain that fire not only when we perform a particular action but also when we watch someone else perform that same action.

Neuroscientists believe this "mirroring" is the mechanism by which we can "read" the minds of others and empathize with them. It's how we "feel" someone's pain, how we discern a grimace from a grin, a smirk from a smile.

Problem was, there was no proof that mirror neurons existed — only suspicion and indirect evidence. Now, reporting in the April edition of the journal <u>Current Biology</u>, Dr. Itzhak Fried, a UCLA professor of <u>neurosurgery</u> and of psychiatry and biobehavioral sciences, Roy



Mukamel, a postdoctoral fellow in Fried's lab, and their colleagues have for the first time made a direct recording of mirror neurons in the human brain.

The researchers recorded both single cells and multiple-cell activity, not only in motor regions of the brain where mirror neurons were thought to exist but also in regions involved in vision and in memory.

Further, they showed that specific subsets of mirror cells increased their activity during the execution of an action but decreased their activity when an action was only being observed.

"We hypothesize that the decreased activity from the cells when observing an action may be to inhibit the observer from automatically performing that same action," said Mukamel, the study's lead author. "Furthermore, this subset of mirror neurons may help us distinguish the actions of other people from our own actions."

The researchers drew their data directly from the brains of 21 patients who were being treated at Ronald Reagan UCLA Medical Center for intractable epilepsy. The patients had been implanted with intracranial depth electrodes to identify seizure foci for potential surgical treatment. <u>Electrode</u> location was based solely on clinical criteria; the researchers, with the patients' consent, used the same electrodes to "piggyback" their research.

The experiment included three parts: facial expressions, grasping and a control experiment. Activity from a total of 1,177 neurons in the 21 patients was recorded as the patients both observed and performed grasping actions and facial gestures. In the observation phase, the patients observed various actions presented on a laptop computer. In the activity phase, the subjects were asked to perform an action based on a visually presented word. In the control task, the same words were



presented and the patients were instructed not to execute the action.

The researchers found that the neurons fired or showed their greatest activity both when the individual performed a task and when they observed a task. The mirror neurons making the responses were located in the medial frontal cortex and medial temporal cortex, two neural systems where mirroring responses at the single-cell level had not been previously recorded, not even in monkeys.

This new finding demonstrates that mirror neurons are located in more areas of the human brain than previously thought. Given that different brain areas implement different functions — in this case, the medial frontal cortex for movement selection and the medial temporal cortex for memory — the finding also suggests that mirror neurons provide a complex and rich mirroring of the actions of other people.

Because mirror neurons fire both when an individual performs an action and when one watches another individual perform that same action, it's thought this "mirroring" is the neural mechanism by which the actions, intentions and emotions of other people can be automatically understood.

"The study suggests that the distribution of these unique cells linking the activity of the self with that of others is wider than previously believed," said Fried, the study's senior author and director of the UCLA Epilepsy Surgery Program.

"It's also suspected that dysfunction of these mirror cells might be involved in disorders such as autism, where the clinical signs can include difficulties with verbal and nonverbal communication, imitation and having empathy for others," Mukamel said. "So gaining a better understanding of the mirror neuron system might help devise strategies for treatment of this disorder."



Provided by University of California - Los Angeles

Citation: Researchers make first direct recording of mirror neurons in human brain (2010, April 12) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2010-04-mirror-neurons-human-brain.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.