

## OB/GYN offices may offer ideal venue for improving vaccine rates among women

April 20 2010

---

Obstetrician/gynecologist offices may be the ideal venue for boosting vaccination rates among women, say researchers at Duke University Medical Center. They reported today on a successful pilot program focused on providing HPV (human papillomavirus) and Tdap (tetanus, diphtheria, pertussis) vaccines to non-pregnant and post-partum women.

The researchers say the program, funded by the U.S. Centers for Disease Control, could be a model for ob/gyn clinics across the country to increase much-needed immunizations among eligible adults.

"Ob/gyns don't typically think of themselves as vaccinators," says Geeta Swamy, MD, Director of Obstetrics Clinical Research at Duke, who presented the findings today at the CDC National Immunization Conference in Atlanta. "Even though we vaccinate pregnant women against a variety of diseases that are screened for during pregnancy, we still tend to think of vaccinations as happening at the offices of pediatricians, primary care physicians and family practitioners. But many women seek medical care from their gynecologists, even after they have children. Their annual gynecologist visit is a good opportunity to discuss preventive care which includes vaccinations."

The North Carolina pilot program was set up to improve HPV vaccination rates among non-pregnant women. Preliminary data from one clinic shows that non-pregnant women were already being offered HPV, but when post-partum women were offered the vaccine, the rate of vaccination jumped from 0 to 44 percent. "These women would not have

been vaccinated if this program was not in place," Swamy said.

Even more significant was the increase in women who received the Tdap vaccine. "Nearly 600 women received the vaccine of the 1000 who were offered it," she said. None had been offered it before.

Reaching women who had not been vaccinated is important because rates of pertussis have been on the rise for the last five years despite CDC recommendations that adults and adolescents receive the newer, single dose if they had not received a tetanus shot within the past two years.

Although the disease is not as serious in adolescents and adults, it is life-threatening among infants who aren't fully immunized until at least one year of age. A recent CDC report found mothers were the primary source of infection in 32 percent of infant pertussis cases. "Ideally we aim to vaccinate [women](#) before they conceive, but any post partum woman should get the vaccine if their last tetanus shot was two years ago or more," says Swamy. "If we can vaccinate new moms, we can provide a cocooning effect that protects their infants from this deadly disease."

Swamy believes the program could easily be implemented in ob/gyn offices nationwide as more emphasis is being placed on [preventive care](#). "We should be thinking of preventive medicine as something that's done at almost any medical encounter, and not just limiting it to the primary care field. It should be provided by all doctors where it's feasible and within the scope of their care."

Provided by Duke University Medical Center

Citation: OB/GYN offices may offer ideal venue for improving vaccine rates among women (2010, April 20) retrieved 2 May 2024 from <https://medicalxpress.com/news/2010-04-obgyn->

[offices-ideal-venue-vaccine.html](https://www.medicalpress.com/offices-ideal-venue-vaccine.html)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.