

# Low Vitamin D Levels Are Related to MS Brain Atrophy, Cognitive Function, Studies Show

April 28 2010

---

(PhysOrg.com) -- Low vitamin D levels may be associated with more advanced physical disability and cognitive impairment in persons with multiple sclerosis, studies conducted by neurologists at the University at Buffalo have shown.

Low vitamin D levels may be associated with more advanced physical disability and [cognitive impairment](#) in persons with multiple sclerosis, studies conducted by neurologists at the University at Buffalo have shown.

Their results, reported at the American Academy of Neurology meeting, held earlier this month, indicated that:

- The majority of MS patients and healthy controls had insufficient vitamin D levels.
- Clinical evaluation and [magnetic resonance imaging](#) (MRI) images show low blood levels of total vitamin D and certain active vitamin D byproducts are associated with increased disability, brain atrophy and [brain lesion](#) load in MS patients.
- A potential association exists between cognitive impairment in MS patients and low vitamin D levels.

The MRI study involved 236 MS patients -- 208 diagnosed with the relapsing-remitting type and 28 with secondary progressive, a more destructive form of MS -- and 22 persons without MS.

All participants provided blood serum samples, which were analyzed for total vitamin D (D2 and D3) levels as well as levels of active vitamin D byproducts. MRI scans performed within three months of blood sampling were available for 163 of the MS patients.

Results showed that only seven percent of persons with secondary-progressive MS showed sufficient vitamin D, compared to 18.3 percent of patients with the less severe relapsing-remitting type.

Higher levels of vitamin D3 and [vitamin D3](#) metabolism byproducts (analyzed as a ratio) also were associated with better scores on disability tests, results showed, and with less brain atrophy and fewer lesions on MRI scans.

Bianca Weinstock-Guttman, MD, UB associate professor of neurology/Jacobs Neurological Institute and director of the [Baird Multiple Sclerosis](#) Center, is first author on the study. Commenting on these results, Weinstock-Guttman said: "Clinical studies are necessary to assess vitamin D supplementation and the underlying mechanism that contributes to MS disease progression."

While lower-than-normal vitamin D status is known to be associated with a higher risk of developing MS, little is known about its relationship to cognitive impairment.

Sarah A. Morrow, MD, UB assistant research professor of neurology/Jacobs Neurological Institute and lead author on the cognitive-impairment study, compared vitamin D levels in blood samples of 136 MS patients with the results of their neuropsychological assessments that

tested multiple types of cognition affected by MS.

"Results showed that MS patients who were impaired on tests of executive function --critical reasoning and abstract thinking -- and the ability to plan and organize, were more likely to be deficient in vitamin D," said Morrow.

"This relationship held true when controlling for the season during which vitamin D was measured, as well as depression, which is known to be associated with lower vitamin D levels." Morrow noted there also was a suggestion that verbal fluency (word generation) and visual-spatial memory (learning and memory of shapes and figures) is more likely to be affected when [vitamin D](#) levels are not sufficient.

Morrow is continuing her research to clarify these relationships.

Provided by University at Buffalo

Citation: Low Vitamin D Levels Are Related to MS Brain Atrophy, Cognitive Function, Studies Show (2010, April 28) retrieved 20 March 2024 from <https://medicalxpress.com/news/2010-04-vitamin-d-ms-brain-atrophy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--