

Children with severe asthma at increased risk of developing COPD

May 16 2010

Children with severe asthma have more than 30 times the risk of developing adult chronic obstructive lung disease (COPD) as adults compared to children without asthma, according to a prospective longitudinal cohort study from the Royal Children's Hospital in Melbourne.

The results will be presented at the ATS 2010 International Conference in New Orleans.

"There is important [epidemiological evidence](#) to suggest that events in childhood that influence lung growth constitute a significant risk for COPD," explained lead author, Andrew Tai, MBBS, FRACP. "The aim of this study was to describe the association between the pattern of childhood [asthma](#) and the risk of developing adult COPD in a longitudinal cohort."

Subjects of the Melbourne Asthma Study were recruited at the age of seven, from a 1957 birth cohort and were assessed regularly until the age of 50. At recruitment, subjects were classified as having no history of wheeze, intermittent asthma (such as viral-induced wheezing), persistent asthma (in the absence of illness), or severe asthma. Of the surviving members of the original group, 197 answered a detailed questionnaire and underwent [lung function](#) testing for the current study.

Subjects who were classified as having severe asthma in childhood had an adjusted risk of COPD of 31.9 times that of children without asthma.

Interestingly, children with mild asthma were not at increased risk of developing adult obstructive lung disease.

"At this stage, children with mild asthma are those who have symptoms of wheeze which are triggered primarily by respiratory infections. A majority [of children with mild asthma] remit by adolescence or adulthood," explained Dr. Tai. "However, children with more severe asthma features tend to have predisposing risk factors (like atopy) and continue to have symptoms of wheeze well into adult life."

"It is important to emphasize that the lung function decline in this group is not increased compared to those with mild or no asthma, as has been raised in some other studies," Dr. Tai continued. "However, lung function in children with severe asthma are reduced in childhood years and decline in adult life to levels consistent with adult obstructive lung disease. Fundamentally, we believe that this severe asthma group start with a lesser baseline lung function and gradually deteriorate to the levels consistent with a diagnosis of COPD. At this stage, there is no data on when airway remodeling occurs in children and hence, its impact on lung function, but there is an emerging relationship between childhood severe asthma and adult obstructive lung disease."

Importantly, this study was performed on a group of children recruited in the 1960s when anti-inflammatory treatment was not available. Studies to date suggest that anti-inflammatory medications do not alter the natural progression of mild childhood asthma, but there are no studies performed in those children with severe asthma.

"There should be greater emphasis on the surveillance and treatment of children with asthma, therefore potentially preventing the development of adult obstructive lung disease," said Dr. Tai.

Researchers still do not fully understand the mechanisms that link severe

childhood asthma with adult COPD, but these findings suggest that appropriate treatment strategies (and surveillance) should be instigated early in life to potentially minimize future risk.

"Early treatment to prevent airway remodeling in childhood may reduce the incidence of this long-term complication of childhood asthma," concluded Dr. Tai. "Currently, there are more than 30 birth cohort studies of varying duration being conducted around the world. In particular, the long-term follow-up of the Tucson birth cohort into young adulthood has shown trends similar to our findings of airway obstruction originating from early life. Clearly, more research to understand the mechanisms and timing of changes within the airway wall, inflammation and function needs to be conducted, applying preferably non-invasive methods in determining potential contributing factors."

Provided by American Thoracic Society

Citation: Children with severe asthma at increased risk of developing COPD (2010, May 16)
retrieved 5 May 2024 from

<https://medicalxpress.com/news/2010-05-children-severe-asthma-copd.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--