

Brief treatment can ease depression in mothers

May 1 2010

When pediatricians see a child for a check-up, they may want to spend a few minutes checking out the mother's well-being, too.

New research suggests that depression is common in disadvantaged mothers, well beyond the postpartum period. In addition, if depressed moms are treated briefly, their symptoms improve as do their perceptions of their child's behavior or the behavior problems themselves.

"Depression in underserved women of childbearing age is extremely common, and nearly one in five mothers of children older than 1 year of age reports moderate to severe [depressive symptoms](#)," said Carol Weitzman, MD, who will present the study findings Saturday, May 1 at the Pediatric Academic Societies (PAS) annual meeting in Vancouver, British Columbia, Canada.

[Maternal depression](#) has been associated with health and development problems in children. Therefore, to optimize children's health, mothers who are depressed should be identified and treated. As parents routinely bring their children for check-ups, this may present an opportunity to identify depressed mothers as so many women do not seek treatment for their symptoms.

Dr. Weitzman and her colleagues at Yale University School of Medicine asked 931 mothers to complete a 16-item measure of depression severity before a well-child visit in a clinic that cares for disadvantaged children.

Women who screened positive were interviewed to confirm that they had depressive symptoms.

Seventy-one mothers with depression were randomized to receive six sessions of on-site [cognitive behavior therapy](#) (CBT) or case management, which consisted of speaking with a social worker and getting assistance with referrals. Their children's social-emotional functioning was measured before and after treatment using the Brief Infant Toddler Social Emotional Assessment (for children younger than 4 years) or the Pediatric Symptom Checklist.

Results showed that 45 percent of mothers screened positive for depressive symptoms (26 percent had mild symptoms, 13 had moderate symptoms and 6 percent had severe symptoms).

"This finding reinforces that depression in mothers is not restricted to the postpartum period, and in fact after the [postpartum period](#) as children get older, the prevalence of maternal depression may be higher," Dr. Weitzman said.

All of the women who were treated with case management or CBT showed improvement in their depression symptoms. The scores improved significantly in the group who received CBT, whereas the case management group did not show a dramatic change.

After treatment with CBT, women who had children younger than 4 also rated them as having fewer behavioral challenges. No significant changes in behavior were reported among the women who received case management or women with children older than 4 in either treatment group.

"The effect of depression in a parent on a child can be significant," Dr. Weitzman said. "Brief on-site treatment can help reduce symptoms of

depression in a mother and may also improve her perspective about her child's behavior."

Provided by American Academy of Pediatrics

Citation: Brief treatment can ease depression in mothers (2010, May 1) retrieved 19 April 2024 from <https://medicalxpress.com/news/2010-05-treatment-ease-depression-mothers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.