

Bariatric surgery in diabetic adults improves insulin sensitivity better than diet

June 21 2010

Gastric bypass surgery improves Type 2 diabetes by other mechanisms in addition to weight loss and does so better than a low-calorie diet despite achieving equal weight loss, a new study finds. The results will be presented Monday at The Endocrine Society's 92nd Annual Meeting in San Diego.

"Our study shows that in the short term, [weight loss](#) by [diet](#) alone does not achieve the same improvements in diabetes as gastric bypass surgery," said the presenting author, Judith Korner, MD, PhD, assistant professor of medicine at Columbia University College of Physicians and Surgeons, New York.

Korner and her colleagues found that gastric bypass surgery better improved insulin sensitivity, the body's ability to successfully clear glucose sugar from the bloodstream into the cells. Insulin sensitivity is impaired in people with Type 2 diabetes, and obesity adds to this problem. The result is a buildup of sugar in the blood.

The study compared the effects on diabetic adults of a low-calorie diet versus Roux-en-Y gastric bypass, the most common gastric bypass procedure. Roux-en-Y gastric bypass decreases the size of the stomach and reroutes the digestive tract to bypass most of the stomach and part of the small intestine. After gastric bypass, many diabetic patients achieve normal blood glucose control or vastly improved control, and some may no longer require diabetes medications.

In the study, seven obese patients with Type 2 diabetes received a daily 800-calorie liquid diet and no surgery, while seven other obese diabetic adults underwent [gastric bypass surgery](#). The study ended when both groups lost the same amount of weight: an average of 8 percent of body weight. However, the surgery-treated patients lost the weight faster: in about 3.5 weeks compared with 8 weeks for the dieters.

Surgical patients were able to discontinue all of their diabetes medications by the study's end, but the dieters reduced their medication use by 55 percent, Korner reported.

The researchers found significant improvements in the surgery group in measures of insulin sensitivity and function of beta cells, the insulin-producing cells in the pancreas. Improvements in [insulin sensitivity](#) in the low-calorie diet group was not statistically significant and beta cell function improved to a lesser extent.

Korner speculated that hormonal changes may be responsible for the improvements resulting from Roux-en-Y surgery in individuals with [Type 2 diabetes](#).

"It will be important to understand how surgery works to produce these results so that we can develop medical therapies of equivalent efficacy," she said.

Provided by The Endocrine Society

Citation: Bariatric surgery in diabetic adults improves insulin sensitivity better than diet (2010, June 21) retrieved 28 April 2024 from <https://medicalxpress.com/news/2010-06-bariatric-surgery-diabetic-adults-insulin.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.