

Flame retardant linked to altered thyroid hormone levels during pregnancy

June 21 2010

Pregnant women with higher blood levels of a common flame retardant had altered thyroid hormone levels, a result that could have implications for fetal health, according to a new study led by researchers at the University of California, Berkeley.

"This is the first study with a sufficient sample size to evaluate the association between PBDE flame retardants and [thyroid function](#) in pregnant women," said the study's lead author, Jonathan Chevrier, a UC Berkeley researcher in epidemiology and in environmental health sciences. "Normal maternal thyroid hormone levels are essential for normal [fetal growth](#) and [brain development](#), so our findings could have significant public health implications. These results suggest that a closer examination between PBDEs and these outcomes is needed."

PBDEs, or polybrominated diphenyl ethers, are a class of organobromine compounds found in common household items such as carpets, textiles, foam furnishings, electronics and plastics. U.S. fire safety standards implemented in the 1970s led to increased use of PBDEs, which can leach out into the environment and accumulate in human [fat cells](#).

Studies suggest that PBDEs can be found in the blood of up to 97 percent of U.S. residents, and at levels 20 times higher than those of people in Europe. Because of California's flammability laws, residents in this state have some of the highest exposures to PBDEs in the world.

"Despite the prevalence of these flame retardants, there are few studies

that have examined their impact on human health," said the study's principal investigator, Brenda Eskenazi, UC Berkeley professor of epidemiology and of maternal and child health. "Our results suggest that exposure to PBDE flame retardants may have unanticipated human health risks."

The new study, to be published June 21 in the journal [Environmental Health Perspectives](#), is the second study to come out this year from Eskenazi's research group linking PBDEs to human health effects. Eskenazi was the principal investigator on the earlier study that found that women with higher exposures to flame retardants took longer to get pregnant.

In the new study, the researchers analyzed blood samples from 270 women taken around the end of their second trimester of pregnancy. The women in the study were part of a larger longitudinal study from the Center for the Health Assessment of Mothers and Children of Salinas (CHAMACOS) that examines environmental exposures and reproductive health.

The researchers measured concentrations of 10 PBDE chemicals, two types of thyroxine (T4) and thyroid-stimulating hormone (TSH). They controlled for such factors as maternal smoking, alcohol and drug use, and exposure to lead and pesticides.

Analysis focused on the five PBDE chemicals that were detected most frequently and are components of a mixture called pentaBDE. The researchers found that a 10-fold increase in each of the PBDE chemicals was associated with decreases in TSH ranging from 10.9 percent to 18.7 percent. When the five PBDEs were analyzed together, a tenfold increase was linked to a 16.8 percent decrease in TSH.

The study did not find a statistically significant effect of PBDE

concentrations on levels of T4. With one exception, all the women in the study with low TSH levels had normal free T4 levels, which corresponds to the definition of subclinical hyperthyroidism. The study found that odds of subclinical hyperthyroidism were increased 1.9 times for each tenfold increase in PBDE concentrations.

"Low TSH and normal T4 levels are an indication of subclinical hyperthyroidism, which is often the first step leading toward clinical hyperthyroidism," said Chevrier. "Though the health effect of subclinical hyperthyroidism during pregnancy is not well understood, maternal clinical hyperthyroidism is linked to altered fetal neurodevelopment, increased risk of miscarriage, premature birth and intrauterine growth retardation."

Exactly how flame retardants influence TSH levels is unclear, the researchers said, but animal studies have shown that certain PBDEs can mimic thyroid hormones.

In addition to the commercial mixture pentaBDE, octaBDE and decaBDE have been developed for use as commercial [flame retardants](#). PentaBDE and octaBDE have both been banned for use by the Stockholm Convention on Persistent Organic Pollutants, the European Union and eight U.S. states, including California, but they are still present in products made before 2004.

The production of decaBDE by major manufacturers is scheduled to be phased out in the United States by 2013. However, pentaBDE and decaBDE are being replaced by new brominated and chlorinated compounds whose impact on human health is not yet clear, the researchers noted.

More information: The study is available for free online: ehponline.org/article/info:doi/10.1289/ehp.1001905

Provided by University of California - Berkeley

Citation: Flame retardant linked to altered thyroid hormone levels during pregnancy (2010, June 21) retrieved 3 April 2024 from

<https://medicalxpress.com/news/2010-06-flame-retardant-linked-thyroid-hormone.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--