

## Study examines outcomes of lowering homocysteine levels with folic acid and vitamin B12

## June 22 2010

Patients who had experienced a heart attack and lowered their blood homocysteine levels with folic acid and vitamin B12 supplementation did not have an associated lower risk of heart attack, coronary death or stroke, according to a study in the June 23/30 issue of *JAMA*. However, the researchers did find that folic acid supplementation did not increase the risk of cancer, which has been speculated.

Blood homocysteine levels are positively associated with <u>cardiovascular</u> disease, but it is uncertain whether the association is causal, according to background information in the article. A meta-analysis of prospective studies indicated that, after adjustment for known risk factors, a 25 percent lower than usual homocysteine concentration was associated with an 11 percent lower risk of coronary <u>heart disease</u> and 19 percent lower risk of stroke. "Daily supplementation with folic acid typically lowers homocysteine levels by about 25 percent, and the addition of vitamin B12 lowers it by a further 7 percent," the authors write. Other research has suggested supplementation with folic acid may offer a protective effect against stroke.

Jane M. Armitage, F.R.C.P., of the University of Oxford, United Kingdom, and colleagues with the Study of the Effectiveness of Additional Reductions In Cholesterol and Homocysteine (SEARCH) trial assessed the effects of lowering homocysteine levels with folic acid plus vitamin B12 in 12,064 survivors of <u>myocardial infarction</u> (heart



attack) in secondary care hospitals in the United Kingdom between 1998 and 2008. Patients were randomized to receive either 2 mg. folic acid plus 1 mg. vitamin B12 daily or matching placebos. The primary outcomes measured included first major vascular event, defined as major coronary event (coronary death, <u>heart attack</u>, or coronary revascularization), fatal or nonfatal stroke, or noncoronary revascularization.

Patients who received the study vitamins reduced homocysteine by an average of  $3.8 \mu mol/L$  (28 percent). During 6.7 years of follow-up, major vascular events occurred in 1,537 of 6,033 participants (25.5 percent) allocated folic acid plus vitamin B12 vs. 1,493 of 6,031 participants (24.8 percent) allocated placebo. "There was no evidence of any benefit beginning to emerge with more prolonged treatment and follow-up," the authors write.

Receipt of study vitamins also was not associated with a significant effect on any stroke (vitamins, 4.5 percent vs. placebo, 4.4 percent); noncoronary revascularizations (vitamins, 3.0 percent vs. placebo, 2.5 percent); or major coronary events (vitamins, 20.4 percent vs. placebo, 19.6 percent). There were no apparent differences in the numbers of deaths attributed to vascular causes or nonvascular causes.

New primary cancers (excluding non-melanoma skin cancer) were diagnosed in 11.2 percent of the participants allocated folic acid plus <u>vitamin B12</u> vs. 10.6 percent allocated placebo, with this difference not being significant.

"Taken together with the previous homocysteine-lowering trials, the results of SEARCH indicate that folic acid supplementation has no significant adverse effects on cancer or other major health outcomes, even if it also produces no beneficial effects on cardiovascular disease. In addition, these results highlight the importance of focusing on drug



treatments (e.g., aspirin, statins, and antihypertensive therapy) and lifestyle changes (in particular, stopping smoking and avoiding excessive weight gain) that are of proven benefit, rather than lowering homocysteine with folic acid-based vitamin supplements, for the prevention of cardiovascular disease," the authors conclude.

More information: JAMA. 2010;303[24]:2486-2494.

Provided by JAMA and Archives Journals

Citation: Study examines outcomes of lowering homocysteine levels with folic acid and vitamin B12 (2010, June 22) retrieved 2 May 2024 from <a href="https://medicalxpress.com/news/2010-06-outcomes-lowering-homocysteine-folic-acid.html">https://medicalxpress.com/news/2010-06-outcomes-lowering-homocysteine-folic-acid.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.