

Study finds a seasonal variance in sleep-disordered breathing in young children

June 7 2010

The prevalence of mild sleep-disordered breathing in young children may fluctuate seasonally, suggests a research abstract that will be presented Monday, June 7, 2010, in San Antonio, Texas, at SLEEP 2010, the 24th annual meeting of the Associated Professional Sleep Societies LLC.

Results indicate that in summer and fall the prevalence of mild sleep-disordered [breathing](#) in elementary-school [children](#) increased steadily from June (21.6 percent) through September (37.2 percent) and then decreased from September through November (6.3 percent). Controlling for potential confounders such as age, [body mass index](#), gender and race showed that the odds of mild sleep-disordered breathing in every month was significantly lower than in September.

"What surprised us most was the dramatic impact that season had on the prevalence of SDB," said principle investigator Edward Bixler, PhD, professor and vice chair for research in the department of psychiatry at Penn State University in Hershey, Pa. "The results are significant because they underscore the importance of evaluating a child's sensitivity to seasonal allergies when diagnosing and treating a child for SDB."

The study involved a random sample of 687 children in grades K-5. Their parents completed a brief questionnaire, and each child was evaluated between June and November during an overnight [sleep](#) study in the sleep laboratory. Mild sleep-disordered breathing was defined as having an apnea-hypopnea index (AHI) of one to five breathing pauses

per hour of sleep.

In a study published in the June 2009 issue of the journal SLEEP, Bixler and his research team reported that nasal problems such as chronic sinusitis and [rhinitis](#) are significant risk factors for mild sleep-disordered breathing in children. However, the extent to which allergies may promote a seasonal variation in sleep-disordered breathing still needs to be determined.

The researchers added that the results may have implications for the development of pharmacologic treatment strategies for mild [sleep-disordered breathing](#) in children.

The American Academy of Sleep Medicine reports that approximately two percent of otherwise healthy young children have obstructive sleep apnea, a common form of SDB that occurs when soft tissue in the back of the throat collapses and blocks the airway during sleep. Most children with OSA have a history of snoring that tends to be loud and may include obvious pauses in breathing and gasps for breath. Parents often notice that the child seems to be working hard to breathe during sleep.

Provided by American Academy of Sleep Medicine

Citation: Study finds a seasonal variance in sleep-disordered breathing in young children (2010, June 7) retrieved 18 April 2024 from <https://medicalxpress.com/news/2010-06-seasonal-variance-sleep-disordered-young-children.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.