

# Low Risk of Traumatic Brain Injury from Roller Coaster Rides, Researcher Says

June 28 2010

---

With July 4th around the bend, if you've ever feared a head injury from a roller coaster ride, it's time to stop worrying and enjoy your local amusement parks. Although a significant body of scientific research has long contended that the physics behind gravitational force isn't enough to cause problems, misconceptions have abounded anyway, said Bryan Pfister, PhD, an assistant professor in the department of biomedical engineering at NJIT.

Pfister recently set out to disprove such [misconceptions](#) in his paper "[Head Motions While Riding Roller Coasters: Implications for Brain Injury](#)," (The American Journal of Forensic Medicine and Pathology, Dec. 2009). The paper makes the case that "there appears to be an extremely low risk of [traumatic brain injury](#) (TBI) due to the head motions induced by [roller coaster](#) rides."

"The risk of TBI while riding roller coasters has received substantial attention. Case reports of TBI around the time of riding roller coasters have led many medical professionals to assert that the high gravitational forces (G-forces) induced by roller coasters pose a significant TBI risk. Head injury research, however, has shown that G-forces alone cannot predict TBI," said Pfister.

"Unfortunately, many professionals have incorrectly used G-force measurements to dramatize the risk of head injury on a roller coaster ride. Indeed, their misinterpretation would also conclude that it is too dangerous to drive a car.

While the exposure to G-forces on roller coasters appears high, it is not more than what you would experience in many everyday activities.”

Pfister used established head injury criteria and procedures to compare the potential of TBI between daily activities and roller coaster riding. Three-dimensional head motions were measured during three different roller coaster rides, a pillow fight, and car crash simulations.

Data was analyzed and compared with published data, using similar analyses of head motions. An 8.05 m/s [car crash](#) led to the largest head injury criterion measure of 28.1 and head impact power of 3.41, over 6 times larger than the roller coaster rides of 4.1 and 0.36. Notably, the linear and rotational components of head acceleration during roller coaster rides were milder than those induced by many common activities.

Pfister, a specialist in neural tissue engineering, received in 2007 a prestigious Faculty Early Career Development Award from the National Science Foundation to support and expand his research into rapid axon stretch growth, a technique for regenerating damaged or diseased nerve cells. He received his doctorate from Johns Hopkins University.

Provided by New Jersey Institute of Technology

Citation: Low Risk of Traumatic Brain Injury from Roller Coaster Rides, Researcher Says (2010, June 28) retrieved 24 April 2024 from <https://medicalxpress.com/news/2010-06-traumatic-brain-injury-roller-coaster.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--