

# Vitamin D deficiency confirmed as common across a range of rheumatic conditions

June 18 2010

---

Two separate studies have shown that vitamin D deficiency is common in patients with a range of rheumatic diseases, with over half of all patients having below the 'normal' healthy levels of vitamin D (48-145 nmol/L) in their bodies. A further study assessing response to vitamin D supplementation found that taking the recommended daily dose did not normalise vitamin D levels in rheumatic disease patients. The results of these three studies were presented today at EULAR 2010, the Annual Congress of the European League Against Rheumatism in Rome, Italy.

A UK study<sup>1</sup> of 180 patients aimed to assess mean levels of vitamin D in patients with inflammatory joint diseases, [osteoarthritis](#) and myalgia ([muscle pain](#) that, when experienced long term may be associated with nutritional deficiency). Data on vitamin D levels were gathered and results showed that 58% of individuals with a rheumatic condition had levels below that clinically considered to be 'sufficient' in healthy subjects (48-145 nmol/L).

An Italian study<sup>2</sup> of 1,191 RA patients aimed to determine a correlation between vitamin D deficiency and several different clinical measures of disease activity. Researchers found that, regardless of supplementation, levels of 25-hydroxyvitamin D (25(OH)D), (a standard clinical measure of vitamin D in the blood), were lower than healthy levels (

Citation: Vitamin D deficiency confirmed as common across a range of rheumatic conditions (2010, June 18) retrieved 1 May 2024 from <https://medicalxpress.com/news/2010-06-vitamin-d->

[deficiency-common-range.html](http://deficiency-common-range.html)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.