

Cancer deaths continue to drop

July 7 2010

The continued drop in overall cancer mortality rates over the last 20 years has averted more than three-quarters of a million (767,000) cancer deaths according to a new report from the American Cancer Society. The American Cancer Society's annual Cancer Statistics article reports that the overall death rate from cancer in the United States in 2007 was 178.4 per 100,000, a relative decrease of 1.3 percent from 2006, when the rate was 180.7 per 100,000, continuing a trend that began in 1991 for men and 1992 for women. In that time, mortality rates have decreased by 21 percent among men and by 12 percent among women, due primarily to declines in smoking, better treatments, and earlier detection of cancer.

The figures come from Cancer Statistics 2010, published early online in *CA: A Cancer Journal for Clinicians*. The report and its consumer version, Cancer Facts & Figures 2010, include the estimated numbers of new cancer cases and deaths for 2010. Society epidemiologists predict there will be 1,529,560 new cancer cases (789,620 in men and 739,940 in women) and 569,490 cancer deaths (299,200 in men and 270,290 in women) in the United States in 2010.

Other highlights of Cancer Statistics 2010 include:

- Cancers of the lung, prostate, and colorectum in men and cancers of the lung, breast, and colorectum in women continue to be the most common fatal cancers. These four cancers account for half of the total cancer deaths among men and women.

- Among men, cancers of the prostate, lung, and colon will account for just over half (52 percent) of all newly diagnosed cancers. [Prostate cancer](#) alone will account for one in four (28 percent, 217,730 cases) of all cancer diagnoses in men. About nine in ten of these new cases of prostate cancer are expected to be diagnosed at local or regional stages, for which five-year relative survival approaches 100 percent.
- The three most commonly diagnosed types of cancer among women in 2010 will be cancers of the breast, lung, and colon, accounting for 52 percent of cancer cases in women. Breast cancer alone is expected to account for 28 percent (207,090) of all new cancer cases among women.
- Cancer incidence rates decreased in men 1.3 percent per year from 2000 to 2006 and in women 0.5 percent per year from 1998 to 2006.
- Death rates for all cancer sites combined decreased 2 percent per year from 2001 to 2006 in males and 1.5 percent per year from 2002 to 2006 in females.
- Mortality rates have continued to decrease across all four major cancer sites in men and in women except for female lung cancer, in which rates have stabilized since 2003 after increasing for several decades.
- Lung cancer incidence rates have been declining in men by 1.8 percent per year since 1991 and appear to be leveling off in women after increasing for many decades.
- Colorectal cancer incidence rates decreased markedly from 1998

through 2006, by 3 percent per year in males and by 2.2 percent per year in females.

- Female breast cancer incidence rates decreased by 2 percent per year from 1999 to 2006. However, close inspection of rates by individual year shows that after a 6 percent decrease from 2002 to 2003 (following publication of the Women's Health Initiative), incidence rates from 2003 to 2006 remained relatively unchanged.
- Among males under age 40 years, leukemia is the most common fatal cancer, while cancer of the lung and bronchus predominates in men aged 40 years and older.
- Among females, leukemia is the leading cause of cancer death before age 20, breast cancer ranks first at ages 20 to 59, and lung cancer is the leading cause of cancer death at ages 60 and older.
- Lung cancer surpassed breast cancer as the leading cause of [cancer death](#) in women in 1987, and is expected to account for 26 percent of all female cancer deaths in 2010
- African American men have a 14 percent higher overall incidence rate and 34 percent higher overall death rate than white men. African American women have a 7 percent lower incidence rate, but a 17 percent higher death rate than white women for all cancer sites combined.
- Among other racial and ethnic groups, cancer incidence and death rates are lower than those in whites and African Americans for all cancer sites combined and for the four most common cancer sites.

- Cancer is the second leading cause of death among children between ages one to 14 years in the U.S., after accidents. The five-year relative survival rate among children for all cancer sites combined improved from 58 percent for patients diagnosed in 1975 to 1977 to 81 percent for those diagnosed in 1999 to 2005.

"This report is yet more proof we are creating a world with more birthdays," said John R. Seffrin, Ph.D., chief executive officer of the American Cancer Society and its advocacy affiliate, the American Cancer Society Cancer Action Network (ACS CAN). "We will build on our progress in the fight against cancer through laws and policies that increase access to cancer prevention, early detection, and treatment services, and with a sustained federal investment in research designed to find breakthroughs in the prevention and treatment of the most deadly forms of cancer."

Each year, Cancer Facts & Figures features a Special Section highlighting a particular topic related to cancer. The Special Section of Cancer Facts & Figures 2010 is "Prostate Cancer." It is estimated that about 1 in 6 men in the U.S. will be diagnosed with prostate cancer during their lifetime and 1 in 36 will die from the disease. Despite the important burden of prostate cancer cases and deaths, and extensive research on its causes, prevention, early detection, and treatment, many uncertainties remain. The purpose of this year's Special Section is to provide information to clinicians, men who are concerned about their risk of prostate cancer, who are making decisions about prostate cancer screening or treatment, or who are undergoing treatment or follow-up, as well as to anyone interested in learning more about prostate cancer.

The annual Cancer Statistics report and its companion consumer publication, Cancer Facts & Figures, have become critical tools for scientists, public health experts, and policymakers in assessing the current burden of cancer. These estimates are some of the most widely

quoted cancer statistics in the world. The Society's leading team of epidemiologic researchers, in collaboration with scientists from the National Center for Health Statistics, compiles and analyzes incidence and mortality data to estimate the number of new cancer cases and deaths for the current year nationwide and in individual states.

The expected numbers of new cancer cases and cancer deaths should be interpreted with caution because these estimates are based on statistical models and may vary considerably from year to year. Not all changes in cancer trends can be captured by modeling techniques and sometimes the model may be too sensitive to recent trends, resulting in over- or under-estimates. For these reasons, the estimates should not be compared from year-to-year to determine trends; age-standardized cancer incidence and death rates are the best way to monitor changes in cancer occurrence and death. Despite these limitations, the American Cancer Society's estimates of the number of new cancer cases and deaths in the current year provide reasonably accurate estimates of the burden of new cancer cases and deaths in the United States. Such estimates will assist in continuing efforts to reduce the public health burden of [cancer](#).

More information: The full report can be viewed at caonline.amcancersoc.org/papbyrecent.dtl

Provided by American Cancer Society

Citation: Cancer deaths continue to drop (2010, July 7) retrieved 17 April 2024 from <https://medicalxpress.com/news/2010-07-cancer-deaths.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.