


# Cells' grouping tactic points to new cancer treatments

July 19 2010

---



This is a cluster of neural crest cells, where the protrusions can be seen at the border of the cluster. Credit: Roberto Mayor, UCL

The study, which used embryonic cells, points to a new way of treating cancer where therapy is targeted at the process of cancer cells grouping together. The aim is to stop cancer cells from spreading and causing secondary tumours.

In order for cells to migrate they form protrusions - much like oars of a boat - in the direction that they want to travel. However, if a single cell is isolated it produces these oars in all directions and ends up rowing in circles. To move around effectively cells must stick together before

attempting to travel.

The study, published today in the journal *Developmental Cell*, explains how this process works. Scientists have discovered that when cells group together the contact with other cells inhibits the formation of protrusions or 'oars'. This means that protrusions only form on the cells that are on the outside edges of the group, causing the group to move in specific direction as the group is pushed by the outermost 'leader' cells.

Dr Roberto Mayor, UCL Department of Cell and Developmental Biology and lead author of the research, said, "Being able to form a group with neighbour cells is advantageous for migration of embryonic cells as well as [cancer](#) cells during tumour [metastasis](#) - they have strength in unity.

"The findings suggest an alternative way in which cancer treatments might work in the future if therapies can be targeted at the process of group formation to stop cancer cells from spreading and causing secondary tumours."

The experiments were carried out using neural crest cells, which are found next to the developing [central nervous system](#) in embryos. These cells can develop into a huge variety of different kind of cells including heart, face, skin and [muscle cells](#). Scientists blocked surface molecules, proteins called N-cadherin, on the neural crest cells that are involved in forming contacts between cells. When they did this the power of the cells to group together was lost, along with any ability to migrate.

It is expected that inhibition of N-cadherin would have the same the effect in [cancer cells](#), which also move in groups during metastasis.

**More information:** 'Collective chemotaxis requires contact dependent cell polarity' is published in the journal *Developmental Cell*.

Provided by University College London

Citation: Cells' grouping tactic points to new cancer treatments (2010, July 19) retrieved 20 April 2024 from <https://medicalxpress.com/news/2010-07-cells-grouping-tactic-cancer-treatments.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.