

Dogs may help collar Chagas disease

July 12 2010

Chagas disease, for example, is caused by a parasite that roams with only limited control among the rural poor in Latin America. The main vector for the parasite Trypanosoma cruzi is the triatomine insect, or "kissing bug," which thrives in the nooks and crannies of mud-brick dwellings. The bug sucks the blood of mammals, helping *T. cruzi* move between wildlife, cats, dogs and humans.

"Dogs tend to lie on porches or other areas easily accessible to the bugs," says disease ecologist Uriel Kitron, chair of environmental studies at Emory University. "And when a dog is malnourished and its immune system isn't great, they are even more at risk."

Kitron has been researching Chagas disease in remote communities of northern Argentina for the past 10 years. "One of our most significant findings is the importance of dogs in both the spread of the disease, and the potential to help control it," he says, explaining that dogs can make good sentinels for health officials monitoring *T. cruzi* transmission.

Chagas disease begins as an acute infection that can subside on its own. In one out of three cases, however, the infection persists and can go unnoticed for decades, until it causes complications such as heart failure, digestive problems and <u>sudden cardiac death</u>. The condition affects 10 to 12 million people in Latin America, killing more than 15,000 a year.

Human migration has moved Chagas disease around the globe: U.S. blood banks must now screen donors for *T. cruzi*. And bugs travel hidden in people's luggage to new places such as Patagonia in southern



Argentina.

Kitron is collaborating with Ricardo Gürtler of the University of Buenos Aires on a research project funded through a joint NIH-NSF program on the ecology of <u>infectious diseases</u>. Their work in Argentina's Chaco province is included in a June 24 special supplement of *Nature*, devoted to the topic of Chagas disease.

"We are interested in answering scientific questions, but we also want to help reduce the risk and the impact of the disease on the rural population," Kitron says.

Few government resources make it to the rural poor, and the main control for Chagas disease is spraying insecticide. "It's a limited strategy," Kitron says. "If you want to control Chagas disease, you have to look at the whole picture."

The researchers have shown, for example, that people with fewer than two dogs in a household are unlikely to become infected. It turns out that dogs are 14 times more effective at spreading <u>Chagas disease</u> than humans.

"Many of the dogs are not in good shape, they're exposed to a whole bunch of parasites and worms and they just get scraps to eat," Kitron says. "But the idea of just eliminating the dogs is not an option. People really care about their dogs."

An alternative may be to identify dogs that are most at risk of remaining infectious for a long period of time. These "super spreaders" could be targeted with insecticide collars. Research is also ongoing for a vaccine against *T. cruzi* in mongrel <u>dogs</u>.



Provided by Emory University

Citation: Dogs may help collar Chagas disease (2010, July 12) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2010-07-dogs-collar-chagas-disease.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.