

HPV DNA testing over 50 percent more sensitive than cytology testing

July 8 2010

HPV DNA testing for precancers and cervical cancers is over 50 per cent more sensitive than cytology testing for the same conditions a community-based study in Mexico has revealed. Results of the largest HPV DNA primary screening study ever to be performed in a Latin American country point to the readiness of HPV DNA testing for large-scale implementation in Mexico and are published online today in *Cancer, Causes and Controls*.

Attila Lorincz, Professor of Molecular Epidemiology at Barts and The London School of Medicine and Dentistry is among researchers behind the study which involved over 50,000 women, aged 20-70, undergoing routine cytology and high risk (HR) HPV DNA testing. All those who tested positive through cytology were referred to colposcopy, while the HPV DNA positive women were followed by repeat HPV DNA testing with colposcopy if they were repeat positive in one year. The study is a pilot implementation to determine the real world performance of routine HPV DNA testing and routine cytology testing in the Mexican IMSS health insurance system.

In women who were both cytology and HPV DNA positive the rate of precancer or [cancer](#) was 19.7 per cent whereas the corresponding rate in women who were positive by cytology alone was only 3 per cent. In women who were positive twice for HR HPV DNA (regarded as HPV persistent) the rate of precancer and/or cancers was 11.7 per cent.

Overall the sensitivity of HPV DNA testing for precancers and cervical

cancers was 93.3 per cent versus only 40 per cent for cytology testing.

Professor Lorincz said of the results: "The data confirms that HR-HPV [DNA testing](#) is much more sensitive than cytology and that HPV testing also identifies women with high risk of serious cervical disease in an efficient and medically acceptable manner.

"The advantage of this new [cervical cancer](#) screening approach is that women who are HPV DNA positive can be quickly referred to colposcopy (within one year), thus catching the vast majority of the precancers and cancers. In contrast women who are HPV DNA negative can be safely made subject to much longer screening intervals (5 to 10 years) saving considerable costs for the Mexican health system. "

More information: 'A pilot study of HPV DNA and cytology testing in 50,159 women in the routine Mexican Social Security Program,' is published online today (8 July 2010) in Cancer Causes & Control.

Provided by Queen Mary, University of London

Citation: HPV DNA testing over 50 percent more sensitive than cytology testing (2010, July 8) retrieved 2 May 2024 from <https://medicalxpress.com/news/2010-07-hpv-dna-percent-sensitive-cytology.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
