

# Study links African ancestry to high-risk breast cancer

July 22 2010

---



Lisa A. Newman, M.D., M.P.H., is with the University of Michigan Health System. Credit: University of Michigan Comprehensive Cancer Center

A new study finds that African ancestry is linked to triple-negative breast cancer, a more aggressive type of cancer that has fewer treatment options.

Researchers at the University of Michigan Comprehensive Cancer Center found that, among women with [breast cancer](#), 82 percent of African women were triple negative, 26 percent of African-Americans were and 16 percent of white Americans were.

Triple negative breast cancer is negative for three specific markers that are used to determine treatment: the estrogen receptor, the progesterone receptor and HER-2/neu.

"The most significant recent advances in breast cancer treatment have involved targeting these three receptors. But these treatments do not help women with triple-negative breast cancer. Outcome disparities are therefore likely to increase, because fewer African-American women are candidates for these newer treatments," says study author Lisa A. Newman, M.D., M.P.H., director of the Breast Care Center at the U-M Comprehensive Cancer Center.

The study, published online in the journal *Cancer*, looked at 581 African American women and 1,008 white women diagnosed with breast cancer at the Henry Ford Health System in Detroit, plus 75 African women diagnosed at the Komfo Anokye Teaching Hospital in Ghana.

Researchers found that Ghanaian women were diagnosed at a younger age than American women, and with larger tumors and more advanced cancer. In addition, the Ghanaian women were more likely to test negative for each of the three markers.

Prior studies have shown that while African-American women are less likely than white women to develop breast cancer, those who are diagnosed are usually younger and are more likely to die from the disease. These characteristics, including the triple negative disease, are also more common among women with a known hereditary predisposition for breast cancer related to BRCA1 gene mutations. Other studies have also shown a hereditary breast cancer risk associated with racial-ethnic identity -- most commonly among Ashkenazi Jewish women.

"[African ancestry](#) might be associated with other links to hereditary predisposition for particular patterns of breast cancer. We hope that by studying breast cancer in African and African-American [women](#) we can identify biomarkers that might be useful for assessing risk or treating triple-negative breast cancer," says Newman, professor of surgery at the

U-M Medical School.

Breast cancer statistics: 194,280 Americans will be diagnosed with breast cancer this year and 40,610 will die from the disease, according to the American Cancer Society

**More information:** Cancer, published online July 13, 2010,  
[DOI:10.1002/cncr.25276](https://doi.org/10.1002/cncr.25276)

Provided by University of Michigan Health System

Citation: Study links African ancestry to high-risk breast cancer (2010, July 22) retrieved 24 April 2024 from  
<https://medicalxpress.com/news/2010-07-links-african-ancestry-high-risk-breast.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.