

Surgery linked to Creutzfeldt-Jakob disease

July 8 2010



Surgery is linked to Creutzfeldt-Jakob disease. Credit: SINC

A new study spearheaded by Spanish scientists demonstrates a causal relationship between the onset of Creutzfeldt-Jakob disease (CJD), caused by a protein called a prion, and general surgery. CJD manifests itself in hereditary acquired; and sporadic forms, or for unknown reasons, which accounts for the majority of cases.

"Based on the monitoring records of spongiform [encephalopathy](#) in two Nordic countries, we studied the possibility of transmission of the sporadic form of CJD through [general surgery](#)", explains Jesús de Pedro, main author of the study and head of prion monitoring in patients at the National Epidemiology Centre of the Carlos III Health Institute.

The finding, published recently in the *Journal of Neurology, Neurosurgery & Psychiatry*, reveals that, with a few exceptions, the risk

of having contracted the sporadic form of CJD manifests itself at least 20 years after having undergone an operation.

"While we are not ruling out the idea that intraoperational transfusions may play a secondary part, the data suggest that the disease enters and spreads much more quickly within the central or peripheral nervous system", confirms De Pedro.

According to the authors, the fact that computer records of surgeries have been in place since the early seventies in hospitals in Sweden and Denmark enables operations on residents of those countries to be linked to cases of CJD, which "extends an extraordinary quality to the information and more credibility to the findings given the almost total absence of memory bias".

Why is the idea of transmission through surgery important?

The most interesting thing about this finding, which points to an external cause that could be prevented, is that "it may signify a shift in our understanding of the nature of neurodegenerative diseases, such as Alzheimer's or Parkinson's".

We might, therefore, ask ourselves if other types of motor neurone diseases can be transmitted through surgery and be latent for decades, such as those where risk factors, particularly physical professions and activities or certain sporting activities, for example, which are more likely to lead to surgery, have already been indicated.

"Suggesting that a disease could have been acquired during health care is a very delicate affirmation, as some relatives of patients with sporadic CJD may be tempted to seek compensation from health authorities for the alleged intraoperational transmission years previously, which would be impossible to prove in individual cases", he reasons.

Nonetheless, the most conclusive pattern that the study presents, albeit based on few cases and one that must be replicated in future studies, is that the onset of CJD occurs approximately 10 years after an operation on the retina with reused equipment.

More information: Jesús de Pedro-Cuesta, Ignacio Mahillo-Fernández, Alberto Rábano, Miguel Calero, Mabel Cruz, Ake Siden, Henning Laursen, Gerhard Falkenhorst, Kare Mølbak y el Grupo de Investigación EUROSURGYCJD. "Nosocomial transmission of sporadic Creutzfeldt-Jakob disease: results from a risk-based assessment of surgical interventions". J Neurol Neurosurg Psychiatry (2010). [doi:10.1136/jnnp.2009.188425](https://doi.org/10.1136/jnnp.2009.188425)

Provided by FECYT - Spanish Foundation for Science and Technology

Citation: Surgery linked to Creutzfeldt-Jakob disease (2010, July 8) retrieved 25 April 2024 from <https://medicalxpress.com/news/2010-07-surgery-linked-creutzfeldt-jakob-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.