


Structural defects precede functional decline in heart muscle

August 30 2010



UI study shows that T-tubule disruption starts to occur even before any decline in heart function is detectable. The study also finds that T-tubule disorganization gradually worsens over the progression of heart disease and correlates with the severity of cardiac hypertrophy and predicts heart function. Understanding how T-tubule disruption occurs may lead to new ways to diagnose or treat heart failure. Credit: University of Iowa

The disruption of a structural component in heart muscle cells, which is associated with heart failure, appears to occur even before heart function starts to decline, according to a new study by researchers at the University of Iowa Roy J. and Lucille A. Carver College of Medicine.

The structure is a highly organized network of grooves in heart muscle membrane called T-tubules. This network is essential for transmitting

electrical signals to the cell's interior where they are translated into contractions that make the heart beat.

It was previously known that T-tubules become very disorganized during [heart failure](#). The new study, published in the Aug. 20 issue of the journal *Circulation Research*, shows that this disorganization starts well before heart failure occurs during a stage known as compensated hypertrophy, when the heart muscle is enlarged but still able to pump a normal amount of blood around the body.

"Although heart function appears normal during compensated [hypertrophy](#), we found that there already is structural damage," said Long-Sheng Song, M.D., senior author of this paper and UI assistant professor of internal medicine. "Our study suggests that things are going wrong very early in the process, and if we could prevent or slow this damage, we might be able to delay the onset of heart failure."

The researchers used a state-of-the-art imaging technique called laser scanning confocal microscope to visualize these structural changes in an [animal model](#) of heart failure. The study compared T-tubule structure and [heart function](#) at different stages of [heart disease](#) and found that the more disorganized the T-tubule network becomes, the worse the heart functions.

Moreover, the researchers found that T-tubule disorganization was also accompanied by a reduction in levels of a molecule called junctophilin-2, which is thought to be involved in formation of T-tubule networks. In cell experiments, loss of this molecule led to reduced T-tubule integrity.

Although the new findings are not ready to be applied in a clinical setting, understanding how T-tubule disruption occurs may lead to new ways to diagnose or treat heart failure.

Provided by University of Iowa

Citation: Structural defects precede functional decline in heart muscle (2010, August 30)
retrieved 5 May 2024 from

<https://medicalxpress.com/news/2010-08-defects-functional-decline-heart-muscle.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--