

10-fold rise in obesity surgery in England since 2000

August 26 2010

The use of bariatric or weight loss surgery has increased ten-fold in NHS hospitals in England since 2000, finds a study published in the British Medical Journal today.

One reason for this rapid rise is increased demand from obese patients as they become more aware of surgery as a viable treatment option, suggest the researchers.

Bariatric surgery is performed on people who are dangerously obese, for the purpose of losing weight. This is usually achieved by reducing the size of the stomach with a surgical band ([gastric banding](#)), by re-routing the small intestines to a small stomach pouch (gastric bypass), or by removing a portion of the stomach (sleeve gastrectomy).

Bariatric surgery has been shown to reduce the risk of death, hospital admission, and long term cost to the health service. It is recommended by the National Institute for Health and Clinical Excellence (NICE) for people with "morbid obesity" ([body mass index](#) of at least 40 or of at least 35 if accompanied by coexisting disease that could be improved by [weight loss](#)) for whom all non-surgical treatments for weight loss have failed. But little is known about who is actually having bariatric surgery in England.

So a team of researchers based at Imperial College London set out to analyse national outcomes after surgery for obesity in the NHS in England.

Using the Hospital Episode Statistics database, they identified all adult patients who had a first elective bariatric procedure (gastric bypass, gastric banding or sleeve gastrectomy) between April 2000 and March 2008.

Mortality rates 30 days and one year after surgery were recorded, as well as duration of hospital stay and unplanned readmission rates.

A total of 6,953 bariatric procedures were carried out during the study period. The number of procedures rose more than ten-fold from 238 in 2000 to 2,543 in 2007.

Patients selected for gastric banding had lower post-surgery mortality and readmission rates and a shorter length of stay than those selected for gastric bypass.

Patients with comorbidities showed poorer post-surgery outcomes than did other patients.

The percentage of laparoscopic (keyhole) procedures also increased during this time, from 28% in 2000 to 75% in 2007. However, no significant increase in mortality or unplanned readmission was seen over the study period, suggesting that laparoscopy has been introduced in a safe manner into the NHS, say the authors.

They suggest: "In conjunction with the growing level of obesity, as patients become more aware of surgery as a viable treatment option, demand for surgery among morbidly obese patients increases."

Provided by British Medical Journal

Citation: 10-fold rise in obesity surgery in England since 2000 (2010, August 26) retrieved 20

March 2024 from <https://medicalxpress.com/news/2010-08-fold-obesity-surgery-england.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.