

# Gallbladder cancer may be linked to estrogens

August 16 2010

---

A very aggressive disease with a poor prognosis, gallbladder cancer may be connected to higher exposure to estrogens, according to a group of researchers at the University of Houston (UH).

Dr. Jan-Lke Gustafsson, Robert A. Welch Professor in UH's biology and biochemistry department, described his team's findings in a paper titled "Estrogen-dependent gallbladder carcinogenesis in LXR $\beta$ -/- female mice" appearing in a recent issue of the journal *Proceedings of the National Academy of Sciences*, one of the world's most-cited multidisciplinary scientific serials.

"For the first time, we show in this paper that the absence of liver X beta receptors, or LXR $\beta$ , in a complex interplay with estrogens, induces gallbladder cancer exclusively in female mice," Gustafsson said.

"Interestingly, the elimination of estrogens prevents the development of tumors in this animal model."

In the study, the team found that chronic inflammatory gall bladder disease characteristic of LXR $\beta$ -/- mice, led to gallbladder lesions that developed and evolved into cancer in older female mice. It is known that metabolic and hormonal alterations have been associated with this invasive disease, and LXR $\beta$  is a sensor for cholesterol derivatives. By removing the ovaries and reducing estrogen levels, the researchers were able to prevent the development of tumors in LXR $\beta$ -/- mice.

There are many crucial clinical implications resulting from these

findings. First, drugs that decrease the level of estrogens might be added to the conventional treatment of gallbladder cancer. And, in the long term, pharmacological activators of LXR $\beta$  could become potential new anti-cancer drugs that may reduce or regulate the proliferation of gallbladder cells.

Additionally, in looking at families affected by hereditary gallbladder cancers, this research could shed light upon mutations in the sequence of LXR $\beta$  that may be responsible for this particular cancer, indicating a higher risk for this disease. This, in turn, could one day be used to determine individual risks.

"Going forward, we need to estimate exactly the levels of LXR $\beta$  and its activators in human gallbladder cancers, particularly in female patients," Gustafsson said. "Once the presence and the function of LXR $\beta$  in the human gallbladder are clear, we are going to test the potential effects of LXR $\beta$  molecules on human gallbladder cancer cells."

In addition to Gustafsson, the UH team on this project consists of post-doctoral students Drs. Chiara Gabbi, who made the discovery, Hyun-Jin Kim and Rodrigo Barros, as well as biology and biochemistry professor Margaret Warner, who has worked with the Gustafsson group since 1986. Marion Korach-Andre' from the Center for Biosciences at the Karolinska Institute in Sweden also contributed to this study.

Provided by University of Houston

Citation: Gallbladder cancer may be linked to estrogens (2010, August 16) retrieved 18 April 2024 from <https://medicalxpress.com/news/2010-08-gallbladder-cancer-linked-estrogens.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.