

Heart's sounds can help diagnose heart failure

August 10 2010

For emergency department patients with shortness of breath and a risk of heart failure, physicians usually grab one thing first: a stethoscope.

It allows them to hear the S3, an abnormal third sound in the heart's rhythm strongly associated with cardiac disease and heart failure. However, the low-frequency, low-pitch sound is notoriously very difficult to hear with a stethoscope alone.

In a study available online in the *American Journal of Emergency Medicine*, a UC emergency medicine physician concludes that acoustic cardiography, a new technology combining a 12-lead ECG with cardiac acoustic data, can aid physicians in detecting the S3—ultimately increasing the accurate diagnosis of acute heart failure in certain subsets of patients.

The study involved analyzing data from one of the largest emergency department-based trials in acute heart failure, the HEart failure and Audicor technology for Rapid Diagnosis and Initial Treatment (HEARD-IT) multinational trial. The trial, conducted at nine sites from March to October 2006, measured the diagnostic accuracy provided by adding acoustic cardiography to an emergency medicine physician's tools.

"The S3 is highly associated with heart failure," says Sean Collins, MD, UC emergency medicine associate professor and lead author of the study. "So we studied how measuring the presence of the S3 changed physicians' impressions of what was going on, how it potentially changed

their workup and treatment for patients."

In this publication, Collins and his colleagues conducted a secondary analysis on the HEARD-IT results, focusing on specific subgroups of patients presenting a harder diagnostic puzzle. Those include obese patients, patients with [kidney failure](#) and patients with an intermediate level of b-type natriuretic peptide, a biomarker which has been associated with acute heart failure and [cardiovascular risk](#).

They found that adding acoustic cardiography to those patients with indeterminate BNP levels improved diagnostic accuracy 22 percent.

"Our findings suggest we diagnose heart failure only about half the time in these patients without acoustic cardiography," says Collins. "With it, we improve the accuracy to about 70 percent. It's pretty helpful in this subset of patients."

Diagnosing them in the emergency department can put patients on the right pathway to treatment more quickly, says Collins, enabling the physician to streamline further testing and treatment. That's a particular concern with heart failure patients, he says, because many of these patients have associated conditions, such as chronic obstructive pulmonary disease, that make pinpointing the cause of their emergency department visit more difficult.

While Collins says that the HEARD-IT trial didn't make the argument for acoustic cardiography to be used in every patient suspected of having [heart failure](#), this secondary analysis suggests it could be helpful in treating these subgroups.

"It shows that we need to get better at listening for the S3," says Collins, "and this study would suggest that if we don't hear it with our ears, that technology like this might be useful."

Provided by University of Cincinnati Academic Health Center

Citation: Heart's sounds can help diagnose heart failure (2010, August 10) retrieved 28 April 2024 from <https://medicalxpress.com/news/2010-08-heart-failure.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.