

## Heart transplant patients with common disorder have high survival rates

August 24 2010

---

Transplant surgery to correct the most common type of genetic heart disease yields similar short-term and potentially greater long-term survival rates as transplant surgery for other heart diseases, according to research reported in *Circulation: Heart Failure*, an American Heart Association journal.

Researchers found similar survival rates one year after heart transplant surgery between hypertrophic cardiomyopathy (HCM) patients (85 percent) and those with other kinds of heart disease (82 percent). Five years post-surgery, survival rates began to diverge with 75 percent of HCM and 70 percent of other patients surviving. At the 10-year mark, survival rates in both groups dropped, although they remained significantly higher in the HCM patients (61 percent) than in those with other heart diseases (49 percent).

"Patients with this disease who are undergoing transplant can expect reasonable long-term [survival rates](#)," said Martin S. Maron, M.D., lead author and assistant professor of medicine, director of the [Hypertrophic Cardiomyopathy](#) Center, and co-director of Advanced [Cardiac Imaging](#) at Tufts Medical Center in Boston, Mass. "That's a crucial clinical message for this small but important subgroup of patients."

Inflammation of the heart muscle (cardiomyopathy) is a serious, potentially fatal disease that can prevent the heart from pumping blood effectively. In HCM - the second most common form of heart muscle disease - the pumping chamber of the heart, known as the [left ventricle](#),

thickens (hypertrophies), making it stiff and less able to relax and for blood to fill the heart chambers.

Investigators used the United Network of Organ Sharing Registry, a nationwide database of all U.S. transplant patients, to analyze 26,706 adult patients' clinical and survival characteristics. HCM patients comprise about 1 percent of all U.S. heart transplant cases. Yet, the survival rate is comparable to surgeries for other reasons.

Study participants were mostly white (81 percent) and male (79 percent), average age 52. HCM patients, however, tended to be younger, average age 43, and more than half were women. Nearly one-third (31 percent) of participants reported smoking, although this rate was much lower among those with HCM (17 percent). All had received a heart transplant between January 1990 and December 2004.

Provided by American Heart Association

Citation: Heart transplant patients with common disorder have high survival rates (2010, August 24) retrieved 18 April 2024 from <https://medicalxpress.com/news/2010-08-heart-transplant-patients-common-disorder.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.