

Not all kidney patients treated equal: Differences in care for patients beginning on dialysis

August 9 2010

Kidney disease patients insured by some federally sponsored national health care organizations are more likely to undergo an important predialysis procedure than patients with other types of insurance, according to a study appearing in an upcoming issue of the *Journal of the American Society Nephrology* (JASN). The results may provide insights into ways to improve kidney-related care for patients before they go on dialysis.

For patients whose kidneys no longer work, hemodialysis performs the work of the kidneys. Experts recommend creating an arteriovenous fistula (AVF), to connect a vein and artery and allow access to the vascular system. An AVF provides a long-lasting site through which blood can be removed and returned during the dialysis procedure, which patients must undergo 3-4 times per week.

While AVF use for hemodialysis access is safe, cost-effective, and associated with improved survival in [kidney disease](#) patients, only a minority of US patients start hemodialysis with an AVF (13.7% in 2008).

Frank Hurst, MD (Walter Reed Army Medical Center and F. Edward Hebert School of Medicine) and his colleagues evaluated the use of AVFs in federally sponsored national health care organizations such as the Departments of Veterans Affairs (DVA) and Department of Defense

(DoD). Dr. Hurst and his team searched the United States Renal Data System (USRDS) database to identify patients who initiated hemodialysis between 2005 and 2006. They found that patients who received predialysis kidney care were 10 times as likely to begin dialysis with an AVF compared with patients who did not receive such care. Patients with DVA/DoD insurance were 40% more likely to initiate dialysis with an AVF compared with patients with other types of insurance. Specifically, 27.2% of patients with DVA/DoD insurance initiated hemodialysis with an AVF, compared with 18.5% of patients reporting employer-group insurance, 16.7% of patients with "Other" insurance, 15.6% of patients with Medicare, 13.0% of patients with Medicaid, and 8.2% of patients who reported no insurance. The analysis also revealed that patients who initiated hemodialysis at a DoD facility were twice as likely to use an AVF.

Experts are attempting to increase the use of AVFs through initiatives such as the Fistula First Breakthrough Coalition, whereby the Centers for Medicare & Medicaid Services, End Stage Renal Disease Networks, and the entire kidney community are working together to ensure that every suitable patient will receive the most optimal form of vascular access. Dr. Hurst's findings show that despite these efforts, AVF use is still highly dependent upon a patient's type of health insurance prior to the initiation of dialysis.

"Further study of these federal systems may identify practices that could improve processes of care across healthcare systems to increase the number of patients who initiate hemodialysis with an AVF," the authors wrote. They added that while AVF use is more frequent in DVA/DoD patients, the proportion of these patients who initiate [hemodialysis](#) with an AVF is still far too low.

Study co-authors include Kevin Abbott, MD (Walter Reed Army Medical Center and F. Edward Hebert School of Medicine); Dominic

Raj, MD (George Washington University); Mahesh Krishnan, MD (Davita Clinical Research); Carlos Palant, MD (Washington DC VA Medical Center and George Washington University School of Medicine); Lawrence Agodoa, MD (National Institutes of Health); and Rahul Jindal, D, PhD (George Washington University, Walter Reed Army Medical Center, and Brookdale University Medical Center).

In an accompanying editorial, Steve Schwab, MD and Kennard Brown, PhD (University of Tennessee Health Science Center) expressed dismay that AVF placement before dialysis, which requires treatment by both a nephrologist and a vascular surgeon, is not covered by Medicare, which supports most coverage for dialysis. "We now have a public policy that treats a catastrophic illness such as renal failure but does not act to prevent or ameliorate that catastrophe earlier," they wrote. The authors advocate for amending policy guidelines so that patients can benefit from seeing specialists early in the course of their disease.

More information: "Arteriovenous Fistulas among Incident Hemodialysis Patients in Department of Defense and Veterans Affairs Facilities" ([doi: 10.1681/ASN.2010010025](https://doi.org/10.1681/ASN.2010010025))

"Immature Public Policy for Vascular Access" ([doi: 10.1681/ASN.2010070771](https://doi.org/10.1681/ASN.2010070771))

Provided by American Society of Nephrology

Citation: Not all kidney patients treated equal: Differences in care for patients beginning on dialysis (2010, August 9) retrieved 3 July 2024 from <https://medicalxpress.com/news/2010-08-kidney-patients-equal-differences-dialysis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.