

## Possible alternate therapy for adults with poorly controlled asthma

September 19 2010

---

A drug commonly used for the treatment of chronic obstructive pulmonary disease (COPD) successfully treats adults whose asthma is not well-controlled on low doses of inhaled corticosteroids, reported researchers supported by the National Heart, Lung, and Blood Institute (NHLBI), part of the National Institutes of Health.

"This study's results show that tiotropium bromide might provide an alternative to other [asthma](#) treatments, expanding options available to patients for controlling their asthma," said NHLBI Acting Director Susan B. Shurin, M.D. "The goal in managing asthma is to prevent symptoms so patients can pursue activities to the fullest."

According to the study, adding tiotropium bromide to low doses of inhaled corticosteroids is more effective at controlling asthma than doubling inhaled corticosteroids alone, and as effective as adding the long-acting beta agonist salmeterol. The results were published online today in the [New England Journal of Medicine](#) and presented at the Annual Congress of the European Respiratory Society in Barcelona, Spain.

Increasing inhaled corticosteroids or supplementing them with long-acting beta agonists like salmeterol are the two preferred treatment options available for adults whose asthma is poorly controlled on low doses of inhaled corticosteroids. However, higher doses of corticosteroids do not improve symptoms for all patients and can have significant side effects, while long-acting beta agonists have come under

scrutiny for their risk of worsening asthma symptoms that could result in hospitalization and, rarely, death.

"Tiotropium relaxes smooth muscle in the airways through a different mechanism than beta agonists, and thus may help people who do not respond well to currently recommended treatments," said study lead Stephen Peters, M.D., Ph.D., of Wake Forest University Baptist Medical Center, Winston-Salem, N.C. "Further analysis of the study data will help us better understand which patients respond best to tiotropium. Then we will need to conduct longer-term studies to establish its safety for asthma patients and to determine its effect on the frequency and severity of asthma exacerbations."

Conducted by the NHLBI's Asthma Clinical Research Network, the study compared three treatment methods: doubling the dose of inhaled corticosteroids alone, supplementing a low dose of inhaled corticosteroids with a long-acting beta agonist (salmeterol), and supplementing a low dose of inhaled corticosteroids with a long-acting anticholinergic drug (tiotropium bromide). Anticholinergics block a part of the autonomic nervous system that can cause airway muscles to contract. The study followed 210 adults whose asthma was not well-controlled on low doses of inhaled corticosteroids alone. Participants received each treatment for 14 weeks with two-week breaks in between, for a total of 48 weeks.

Tiotropium bromide was shown to be effective using several asthma control measurements, including patients' day-to-day lung function as well as the number of days in which they had no asthma symptoms and did not need to use their albuterol rescue inhalers. When patients began the trial, their average number of such "asthma control days" was 77 per year (extrapolated from the treatment period). Doubling corticosteroids gave patients another 19 symptom-free days on average, while adding tiotropium to low-dose corticosteroids gave them another 48.

"Much research over the last century has explored the role of cholinergic mechanisms [which constrict the airways] and anticholinergic therapies in asthma. However, this is the first study to explore adding an anticholinergic inhaler to low-dose [inhaled corticosteroids](#)," said James Kiley, Ph.D., director of the NHLBI's Division of Lung Diseases. "The Asthma Clinical Research Network is designed to address exactly these kinds of practical and important management questions, with the ultimate goal of helping asthma patients."

Provided by National Institutes of Health

Citation: Possible alternate therapy for adults with poorly controlled asthma (2010, September 19) retrieved 26 April 2024 from <https://medicalxpress.com/news/2010-09-alternate-therapy-adults-poorly-asthma.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--