

Asthma and cavities both common in kids but not linked

September 16 2010

There is no apparent link between asthma and tooth decay, according to a study published in the September 2010 issue of the *Journal of the American Dental Association*.

"Is There a Relationship between [Asthma](#) and Dental Caries?: A Critical Review of the Literature" examined the 27 separate studies which looked for a link between asthma and [cavities](#) that were reported in 29 papers published between 1976 and March 2010.

"The notion that there is a link between asthma and tooth decay may have its origin in anecdotal statements by emergency room workers who see children with poorly managed asthma. These children could also be more likely to have poorly managed dental conditions, and therefore tooth decay. It's reasonable to believe that poor clinical management may be associated with both conditions, not the asthma that is causing the cavities," said Gerardo Maupomé, B.D.S., M.Sc., Ph.D., professor of preventive and community dentistry at the Indiana University School of Dentistry and a Regenstrief Institute affiliated scientist, who is the first author of the new *JADA* study.

"We found little evidence to suggest that asthma causes tooth decay. In fact, the two largest studies we reviewed found that children with asthma appear to have fewer cavities than others. This may be because their parents are used to taking them to health-care providers, and routinely bring them to the dentist," said Dr. Maupomé.

The large number of variables involved, including severity of asthma symptoms and the variety of types of treatment for the disease, has made it difficult to unequivocally determine whether there is a causal link between the two.

While not apparently associated, tooth decay and asthma are the two most prevalent chronic childhood diseases in the United States.

Routine home and professional dental care are critical for all children. Parents of children with asthma do not need to be concerned about an increased risk of [tooth decay](#) but Dr. Maupomé points out that children who use nebulizers to control their asthma may be inadvertently increasing their frequency of exposure to sugars because these nebulizers use fructose to deliver therapy. The frequency and the amount of certain sugars consumed are major factors leading to cavities.

He also recommends that children who are mouth breathers or who have mouth dryness be checked periodically by their dentists. These conditions may be associated with asthma but they are also found in [children](#) who do not have asthma. Many medications used for the long term (such as asthma medications) have been found to reduce the amount of saliva, which is the first protection of teeth.

Provided by Indiana University School of Medicine

Citation: Asthma and cavities both common in kids but not linked (2010, September 16) retrieved 17 August 2024 from <https://medicalxpress.com/news/2010-09-asthma-cavities-common-kids-linked.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--