

No sponge left behind

September 30 2010



A sponge left inside a patient is visible in the lower right quadrant of this X-ray image. Credit: Image provided by Christopher C. Rupp, M.D.

Using the same technology found in clothing tags used in retail store tracking systems, a study from the University of North Carolina at Chapel Hill shows that surgical sponges with implanted radio frequency (RF) tags may be an effective adjunct to manual counting and X-ray detection in preventing sponges from being left behind in patients following a surgical procedure.

UNC surgeons will report the preliminary results of the study on Tuesday, Oct. 5 during a poster session at the 2010 Annual Clinical Congress of the American College of Surgeons in Washington, D.C.

The reported incidence of a sponge or another foreign body, such as a surgical instrument, being left behind after operations has varied widely



over the years. Previously published reports have estimated ranges of one in 1,000 operations to one in 18,000.

"Our preliminary data agrees with the previously reported incidence of retained surgical sponges," said lead investigator UNC gastrointestinal surgeon Christopher C. Rupp, MD. The researchers used a radiofrequency (RF) detection device in 1,600 operations, and found a sponge in one operation in which manual counting of the sponges was correct.

Sponges used to absorb fluids and improve access to organs during surgical procedures are much different than household sponges. Surgical sponges are mostly made of cotton, and come in sizes of 12-by-12 inches or as small as 4-by-4 inches. These sponges can become difficult to see during an operation because they can mold into different shapes and take on the same color of the fluids being absorbed. Furthermore, the sponges can migrate to other areas of the operative field, and they can be difficult to feel with surgical gloves.

Surgeons use various methods to track sponges during operations. The most common is manual sponge counts by a nurse, but more sophisticated methods include the radio-frequency tagged sponges used in the UNC study, X-ray of the abdomen and bar-coded sponges. The UNC investigators did not include bar-coding systems in their study. Human error involved with manual counting and X-ray interpretation were the impetus that resulted in the studies inception, Rupp reported.

The RF-tagged system in the UNC study has the same technology found in the clothing tags used in retail store tracking systems and in microchips embedded in pets. During surgery, a nurse passes a wand over the patient's body to pick up readings from the RF tags. Newer versions have detection hardware built into the mat the patient lies on. "RF detection is not going to replace counting in the operating room, but it can be used as an adjunct because, from what we're seeing in the



preliminary data, it adds a lot to the safety of the procedure," Rupp said.

"Any foreign body present long enough has a risk of causing infection," he explained, noting there are patients in whom sponges have eroded into other organs, mainly the intestines. "People can come back with chronic pain issues after an operation that also leads to detection of a retained surgical sponge," which may require being hospitalized for additional surgery.

Provided by University of North Carolina School of Medicine

Citation: No sponge left behind (2010, September 30) retrieved 2 May 2024 from <u>https://medicalxpress.com/news/2010-09-sponge-left.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.