

Childhood cancer survivors face long-term risk of GI complications, study finds

October 22 2010

People treated for cancer when they were children have a higher-than-average risk of gastrointestinal problems – some mild, some quite severe – in the years following treatment, according to a study to be presented at the 42nd Congress of the International Society of Paediatric Oncology (SIOP) in Boston on Sunday, Oct. 24. Investigators at Dana-Farber/Children's Hospital Cancer Center and their colleagues say their findings suggest that childhood cancer survivors and their physicians should be especially aware of the increased likelihood of gastrointestinal troubles and factor them into plans for monitoring survivors' health.

"Current treatments have dramatically increased survival rates for children with [cancer](#), but we know that many cancer therapies – including surgery, radiation, and chemotherapy – can cause significant gastrointestinal [GI] complications for patients," says the study's senior author, Lisa Diller, MD, director of the Perini Family Survivors Center at Dana-Farber. "Little is known, however, about the long-term GI consequences for [childhood cancer](#) survivors."

To identify such delayed effects, researchers searched for instances of upper GI (esophageal, stomach) problems, liver disorders, and lower GI (intestinal) problems in 14,358 participants in the Childhood Cancer Survivor Study, which tracks the health of children diagnosed with cancer between 1970 and 1986 who survived at least five years after treatment. Rates of GI disorders in these survivors were compared to rates in a randomly selected group of their siblings.

Most of the survivors had received chemotherapy, and about a third had received radiation and/or chemotherapy. Researchers found that the survivors had an elevated risk for upper GI complications such as ulcers, indigestion, heartburn, esophagus problems, nausea, and vomiting; for liver conditions such as cirrhosis, gallstones, and jaundice (and were 24 percent more likely than siblings to have needed a liver biopsy); and for lower GI problems such as colitis, constipation, diarrhea, and intestinal polyps or blockage. In general, survivors who were diagnosed at an older age and had more intensive radiation and/or chemotherapy and more extensive surgery had higher rates of these complications.

These increased risks make it important for survivors to inform their physicians of their earlier cancer treatment, and for physicians to consider whether GI problems are symptoms of a more serious disease in these patients, Diller states. In recent years, pediatric oncologists have worked to develop lower-dose treatments and less invasive surgical procedures to reduce the long-term consequences of childhood cancer therapy, she adds.

Provided by Dana-Farber Cancer Institute

Citation: Childhood cancer survivors face long-term risk of GI complications, study finds (2010, October 22) retrieved 18 April 2024 from <https://medicalxpress.com/news/2010-10-childhood-cancer-survivors-long-term-gi.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.