

Women with anorexia nervosa more likely to have unplanned pregnancies

October 29 2010

A new study by University of North Carolina at Chapel Hill and Norwegian researchers has found that women with anorexia nervosa are much more likely to have both unplanned pregnancies and induced abortions than women who don't have the serious eating disorder.

These results may be driven by a mistaken belief among [women](#) with anorexia that they can't get pregnant because they are either not having menstrual periods at all or are having irregular periods, said Cynthia M. Bulik, PhD, the study's lead author and director of the UNC Eating Disorders Program.

"Anorexia is not a good contraceptive," Bulik said. "Just because you're not menstruating, or because you're menstruating irregularly, doesn't mean you're not at risk for becoming pregnant."

Physicians and other health care providers need to be aware of this as well, Bulik said. Doctors who treat women and adolescent girls, in particular, "need to make sure that they have the conversation about sexuality and contraception as clearly with patients with anorexia as they do with all other girls and women."

In addition, providers who take care of pregnant women need to know when their patients have an [eating disorder](#) in order to provide appropriate care. Screening for eating disorders during prenatal visits would be an excellent first step, Bulik said.

In the study, published in the November 2010 issue of the journal *Obstetrics & Gynecology*, Bulik and study co-authors analyzed data collected from 62,060 women as part of the Norwegian Mother and Child Cohort Study. There were 62 women in this sample who reported having [anorexia nervosa](#).

The differences between women with anorexia and women with no eating disorder were striking. The average age of the mothers at delivery was 26.2 years in women with anorexia, compared with 29.9 years in the referent group of women without eating disorders. Fifty percent of women with anorexia reported unplanned pregnancies, compared with 18.9 percent, while 24.2 percent of women with [anorexia](#) reported having induced abortions in the past, compared to 14.6 percent.

Provided by University of North Carolina School of Medicine

Citation: Women with anorexia nervosa more likely to have unplanned pregnancies (2010, October 29) retrieved 6 May 2024 from <https://medicalxpress.com/news/2010-10-women-anorexia-nervosa-unplanned-pregnancies.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
