

Combining aerobic and resistance training appears helpful for patients with diabetes

November 23 2010

Performing a combination of aerobic exercise and resistance training was associated with improved glycemic levels among patients with type 2 diabetes, compared to patients who did not exercise, according to a study in the November 24 issue of *JAMA*. The level of improvement was not seen among patients who performed either aerobic exercise or resistance training alone.

Although it is generally accepted that regular exercise provides substantial health benefits for individuals with [type 2 diabetes](#), the exact exercise type (aerobic vs. resistance vs. both) has been unclear. "Given that the 2008 Federal Physical Activity Guidelines recommend aerobic exercise in combination with [resistance training](#), the unanswered question as to whether for a given amount of time the combination of aerobic and resistance exercise is better than either alone has significant clinical and public health importance," the authors write.

Timothy S. Church, M.D., M.P.H., Ph.D., of Louisiana State University System, Baton Rouge, La., and colleagues conducted the HART-D trial, which compared among 262 sedentary women and men with type 2 [diabetes](#) the effect of aerobic training, resistance training, and a combination of both on change in hemoglobin A1c levels (HbA1c; a minor component of hemoglobin [the substance of [red blood cells](#) that carries oxygen to the cells] and to which glucose [blood sugar] is bound; HbA1c levels are used to monitor the control of diabetes mellitus). Study participants were 63.0 percent women, 47.3 percent nonwhite, average age of 56 years, HbA1c level of 7.7 percent and duration of diabetes of

7.1 years. The individuals were enrolled in the 9-month exercise program between April 2007 and August 2009. Forty-one participants were assigned to the nonexercise control group; 73 to resistance training sessions; 72 to aerobic exercise sessions; and 76 to combined aerobic and resistance training.

The researchers found that the absolute change in HbA1c in the combination training group vs. the control group was -0.34 percent. In neither the resistance training (-0.16 percent) nor the aerobic (-0.24 percent) groups were changes in HbA1c significant compared with those in the control group. The prevalence of increases in hypoglycemic medications were 39 percent in the control, 32 percent in the resistance training, 22 percent in the aerobic, and 18 percent in the combination training groups.

"Only the combination exercise group improved maximum oxygen consumption compared with the control group. All exercise groups reduced waist circumference from [-.75 to -1.1 inches] compared with the control group," the authors write. The resistance training group lost an average of 3.1 lbs. fat mass and the combination training group lost an average of 3.7 lbs., compared with the control group.

"The primary finding from this randomized, controlled exercise trial involving individuals with type 2 diabetes is that although both resistance and [aerobic training](#) provide benefits, only the combination of the 2 were associated with reductions in HbA1c levels," the researchers write. "It also is important to appreciate that the follow-up difference in HbA1c between the combination training group and the control group occurred even though the control group had increased its use of diabetes medications while the combination training group decreased its diabetes medication uses."

More information: *JAMA*. 2010;304[20]:2253-2262.

Provided by JAMA and Archives Journals

Citation: Combining aerobic and resistance training appears helpful for patients with diabetes (2010, November 23) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2010-11-combining-aerobic-resistance-patients-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.